ZooKeys 1270: 99–155 (2026)
DOI: 10.3897/zookeys.1270.180704

Research Article

The Swedish Thripidae (Thysanoptera) – with additional notes on wetland thrips

Emma Wahlberg[1], Carl-Axel Gertsson[2], Manfred R. Ulitzka[3], Erika Frisk[1], Ellen Nein[1], Jonas Sandström[4]

1 *Department of Zoology, Swedish Museum of Natural History, P. O. Box 50007, SE-104 05 Stockholm, Sweden*
2 *Murarevägen 13, SE-227 30 Lund, Sweden*
3 *Thrips-iD, Straßburger Straße 37A, 77652 Offenburg, Germany*
4 *SLU Swedish Species Information Centre, Box 7007, SE-750 07, Uppsala, Sweden*

Corresponding author: Emma Wahlberg (emma.wahlberg@nrm.se; emma.wahlberg@arthropoda.se)

Abstract

The Swedish fauna of thrips (Thysanoptera) within the family Thripidae is reviewed in this study. Thripidae is one of the most diverse thrips families, containing many economically important pest species, as well as species of ecological significance. The current investigation updates the knowledge of this family in Sweden, detailing the distribution, ecological associations, and providing the first comprehensive checklist for Swedish Thripidae. We present new records, including first records for Sweden and several regional records, with a detailed morphological identification key. Ten new species are recorded for Sweden, and a total of 98 new provincial records are presented. A total of 567 COI barcode sequences were generated, representing 49 species, and uploaded to BOLD.

Key words: First records, identification, morphology, Sweden, taxonomy, Thripidae, wetlands

OPEN ACCESS

Academic editor:
Elison Fabricio B. Lima
Received: 28 November 2025
Accepted: 9 February 2026
Published: 20 February 2026

ZooBank: https://zoobank.org/DE859176-67D6-4E11-9A38-5A4210F77130

Citation: Wahlberg E, Gertsson C-A, Ulitzka MR, Frisk E, Nein E, Sandström J (2026) The Swedish Thripidae (Thysanoptera) – with additional notes on wetland thrips. ZooKeys 1270: 99–155. https://doi.org/10.3897/zookeys.1270.180704

Copyright: © Emma Wahlberg et al.
This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0).

Introduction

Thrips (Order Thysanoptera) constitute a diverse group of small insects, ranging typically from 0.5 mm to approximately 3 mm in length. The family Thripidae Stephens, 1829, commonly referred to as “common thrips,” is one of the most species-rich groups within the Thysanoptera. Many species within this family have significant agricultural implications, often causing damage to plants by feeding or through the transmission of viruses.

The knowledge of Thripidae in Sweden is based on mainly the larger collections of e.g. Ahlberg (1926) and Qvick (1977). The revision provided by Vasiliu-Oromulu et al. (2001) was based on Ahlberg’s collection revisited. In recent years several papers have proved that thrips offer a lot of new knowledge to discover regarding the Swedish species richness and the distribution of the various species (e.g., Gertsson 2015; Gertsson and Fägerström 2017; Gertsson 2022; Gertsson et al. 2022; Wahlberg 2023b). But the family Thripidae has not previously been comprehensively documented, leaving substantial gaps in our knowledge of species distribution, biology, and ecology. Previous faunistic

studies primarily focused on limited geographical areas or specific pest species. Furthermore, there are no up-to-date identification keys to the Swedish Thripidae. Available keys are older and/or cover other regions (Mound et al. 1976, 2018; Qvick 1977; zur Strassen 2003).

Prior to the material produced herein, the Barcode of Life Database (BOLD, Ratnasingham and Hebert 2007) had only 22 species representing Sweden, with a total of 180 specimens, with molecular COI barcodes. All results are from bycatches from previous projects treating Phlaeothripidae and Aeolothripidae (Wahlberg and Gertsson 2022; Wahlberg 2024). Beside molecular species identification, the COI barcoding sequence can be used to test species concepts with molecular species delimitation methods in combination with morphological examination (Ranasinghe et al. 2022; Emerson 2025). Two species treated here have unclear taxonomic status, *Chirothrips ambulans* Bagnall, 1832 and *Thrips menyanthidis* Bagnall, 1923. *Chirothrips ambulans* was synonymised with *Chirothrips manicatus* by zur Strassen (1959), but later treated as valid species (zur Strassen 2003). However, in e.g. Mound et al. (2018) it is again treated as synonym of *C. manicatus*. As described by Mound et al. (2018), *Thrips menyanthidis* has a likewise unsettled history, synonymised with *Thrips fuscipennis* Haliday, 1836 by Mound et al. (1976) and later validated by zur Strassen (1982). To clarify these taxonomic issues, we have performed species delimitation tests.

This paper aims to consolidate current knowledge and extend the faunistic understanding by compiling an extensive checklist and providing an identification key tailored to Swedish diversity. We also add additional notes on wetland species based on data collected during the research projects on Swedish thrips spanning 2020–2025, involving all families in the region.

Materials and methods

Specimens were collected from various habitats across Sweden, including agricultural fields, gardens, natural grasslands, forests, and urban areas, during 2020–2025. Sampling methods included sweep-netting, beat-tray sampling, malaise traps, and targeted collection from flowers and foliage. Samples have also been donated throughout the project time from both colleagues and the public. A Thysanoptera workshop in August 2024 at the Tovetorp research station also yielded several findings, including new records for Sweden.

Specimen preparation involved maceration and mounting on microscope slides. Specimens collected by E. Wahlberg for DNA extraction were collected and stored in 80% ethanol, prepared according to the non-destructive protocol described in Wahlberg (2023a) and mounted in Euparal. The DNA extraction was carried out with the KingFisher™ Flex (Thermo Scientific) robot and the MagBind® Blood & Tissue DNA HDQ 96 Kit (Omega Bio-tek), according to the manufacturer's protocol. The primer pair LCO1490 and HCO2198 (Folmer et al. 1994) was used for targeting the COI barcoding region, both flanked with a 15 bp oligo in the 5' end to enable subsequent indexing PCR. Amplification was performed with Cytiva Hot Start Mix Ready-To-Go beads and 21 µl H2O, 1 µl of respective primer and 2 µl DNA template. The PCR reaction was run with an initial 95 °C denaturation step (5 min) followed by 38 cycles of 95 °C (30 s), 50 °C (30 s), 72 °C (50 s), finishing with 72 °C (8 min). All PCR products were visualised on a

1% agarose gel and purified using Exonuclease I and FastAP Thermosensitive Alkaline Phosphatase. The indexing PCRs were done in 25 µl reactions, with 0.625 U of Accuprime Pfx DNA Polymerase, 1× AccuPrime reaction mix (Life Technologies, Carlsbad, CA, USA), 0.2 µM of each indexing primer (forward: AATGATACGGCGACCACCGAGATCTACACxxxxxxxxxxACACTCTTTCCCTAC, reverse: CAAGCAGAAGACGGCATACGAGATxxxxxxxxxxGTGACTGGAGTTCAG) and 2 µL of purified PCR product from the first PCR. Cycling conditions were an initial 95 °C (2 min) followed by 8 cycles of 95 °C (15 s), 55 °C (30 s), 38 °C (1 min). The indexing primer design was based on the Adapterama iTru system (Glenn et al. 2019), but with Illumina's 10 bp long Nextera indexes.

All the indexed samples were pooled based on gel band intensity after the first PCR, 8 µl from weak bands and 2 µl from strong ones. The amplicon pool was purified using Cytiva Sera-Mag Select and prepared for sequencing with the Oxford Nanopore Technologies (ONT) LSK-114 ligation kit, and sequenced on a MinION R.10.4.1 FlowCell using MinKNOW 24.06.16 for Linux. Basecalling was set to Super-accurate using Dorado 7.6.8. Barcode splitting was done in MinKNOW using a custom made demultiplex file prepared according to ONTs instructions. During this process, adapter sequences and barcode sequences were trimmed.

Amplicon sorter (Vierstraete and Braeckman 2022) was used to cluster amplicons based on similarity in sequence and length and for building consensus sequences. This method sorts all reads de-novo and reference free, and clusters them into similarity groups. For a greater chance of detecting rare variants, the random sampling option was used, and the --maxreads option was set to a bit higher than double the number of reads in the dataset. Each consensus sequence identified by amplicon sorter was validated in ONT's amplicon workflow in Epi2me (https://github.com/epi2me-labs/wf-amplicon), through Nextflow. (versions: python 3.8.19; fastcat 0.18.6; ezcharts 0.11.2; dominate 2.9.1; numpy 1.24.4; pandas 2.0.3; pysam 0.22.0; si-prefix 1.3.3; seqkit 2.8.2; porechop 0.2.4; samtools 1.19.2; minimap2 2.28-r1209; mosdepth 0.3.7; miniasm 0.3-r179; racon 1.5.0; csvtk 0.27.2; medaka 1.12.0) (Ewels et al. 2020). In this process, the barcode sorted fastq reads were given as input data, and the list of amplicon sorter consensus sequences from the corresponding barcode was passed as reference. In order to investigate the entire read set of each barcode, the options --drop_frac_longest_reads and --reads_downsampling_size were set to 0. Consensus sequences that received more than 5% support, and displayed an even coverage, were further processed. Geneious v. 8.1.6 was used to align the consensus sequences to a reference COI sequence, to determine sequence directionality, and to identify and trim target primer sequences. Each validated variant was exported in fasta format, and finally matched at NCBI Blast (Camacho et al. 2009) and screened for contaminations in NCBI GenBank (Benson et al. 2012). All successfully amplified and sequenced barcodes were uploaded to BOLD and added to the publicly available dataset "Swedish Thysanoptera 2025" (https://dx.doi.org/10.5883/DS-SETHY25) as well as the GenBank sequence database. BOLD and GenBank accession numbers are provided in Suppl. material 1.

Other specimens were collected in AGA (alcohol, glycerine, acetic acid) solution, prepared according to the protocol described in Kobro (2011), and mounted in Canada balm. Additional previously collected material was

examined from the collections at The Swedish Museum of Natural History (**NHRS**) and Lund Museum of Zoology (**MZLU**), and previous distribution data was collected from Kobro (2011), Gertsson (2015), Gertsson and Fägerström (2017), Gertsson (2019), Gertsson (2021), and Wahlberg et al. (2025). Morphological observations were performed using Nikon Eclipse and Swift microscopes equipped with digital imaging systems. Photographs were captured using focus stacking software (Helicon Focus and Swift Imaging), followed by editing in Adobe Photoshop. All material, physical specimens and DNA extract, is deposited at NHRS or Manfred R. Ulitzkas collection with corresponding voucher IDs/collection numbers.

The species concept of *Chirothrips ambulans* and *Thrips menyanthidis* was tested using molecular species delimitation based on the COI barcoding gene sequence. Sequences for available specimens for each species were included, as well as sequences from other species in the same genera. We included sequences from both male and female specimens as well as different geographical locations. The sequences were aligned using MAFFT 7.407 (Katoh and Standley 2013) with auto flavour and standard settings with adjustment of direction. The alignment examined for ambiguous/problematic regions with BMGE 1.12 (Criscuolo and Gribaldo 2010) (sliding window: 3, maximum entropy threshold: 0.5, gap cut of: 0.5, minimum block size: 5). The resulting length of the cleaned alignment was 655 bp. We delimited putative distance-based species with ASAP (Puillandre et al. 2021) with default setting, selecting the best-scoring ASAP partition. For a tree-based delimitation a maximum-likelihood (ML) phylogeny was inferred with IQ-TREE v. 3.0.1 (Nguyen et al. 2015), with ModelFinder model selection (Kalyaanamoorthy et al. 2017), UFBoot2 (1,000 replicates) (Hoang et al. 2018), and SH-like aLRT (1,000 replicates), we applied codon partitioning by position ($1^{st}/2^{nd}/3^{rd}$) with model merging enabled. We applied mPTP 0.2.4 (Kapli et al. 2017) to the phylogeny under the multi-rate model with maximum-likelihood optimisation (--ml --multi). Support was further assessed using MCMC (5,000,000 steps; logging every 10,000; burn-in 100,000).

Remarks on habitats and feeding preferences were collected from Mound et al. (1976, 2018), Schliephake et al. (1979), zur Strassen (2003), original descriptions, and field observations during collection events. Species names follow the current taxonomy as reflected at ThripsWiki (ThripsWiki 2026).

Swedish faunistic provinces and abbreviations

Sweden is traditionally divided into faunistic provinces based on historical cultural regions, which are widely used in entomological literature (Fig. 1). These provinces serve as standardised and stable geographical references and are abbreviated as follows: **Sk** = Skåne, **Bl** = Blekinge, **Ha** = Halland, **Sm** = Småland, **Öl** = Öland, **Go** = Gotland, **GS** = Gotska Sandön, **Ög** = Östergötland, **Vg** = Västergötland, **Bo** = Bohuslän, **Ds** = Dalsland, **Nä** = Närke, **Sö** = Södermanland, **Up** = Uppland, **Vs** = Västmanland, **Vr** = Värmland, **Dr** = Dalarna, **Gä** = Gästrikland, **Hs** = Hälsingland, **Me** = Medelpad, **Hr** = Härjedalen, **Jä** = Jämtland, **Ån** = Ångermanland, **Vb** = Västerbotten, **Nb** = Norrbotten, **Ås** = Åsele lappmark, **Ly** = Lycksele lappmark, **Pi** = Pite lappmark, **Lu** = Lule lappmark, **To** = Torne lappmark.



Figure 1. Map of Sweden with faunistic provinces. Dashed red line represents the Limes Norrlandicus border.

Characters

The identification of Thripidae species is primarily based on microscopic morphological characters (Fig. 2). Key distinguishing features include the shape, number, and position of antennal segments and associated sense cones; presence or absence of ocellar setae; structure and length of wings, wing fringes, and wing veins and setae; presence and arrangement of setae on the pronotum and on the abdominal segments (Fig. 2A). Morphological variations among species and within species at different life stages or populations require careful examination under high magnification.

Results

Taxonomy

Family Thripidae Stephens, 1829

Diagnosis. The family Thripidae can generally be identified by a slender body, the presence of fringed wings, a distinct conical antennal segments often bearing sensory cones, and a saw-like and downwardly turned ovipositor in females (Fig. 2B). Males possess more or less sclerotised genitalia (e.g., phallus, endotheca, and aedeagus), and sometimes pheromone-producing glandular areas on abdominal sternites called pore plates (Fig. 2C). Adult Thripidae typically display narrow wings with marginal fringes, distinctive wing veins, and antennae that frequently exceed the body width.



Figure 2. Morphological characters. **A.** Habitus of *Thrips fuscipennis*, female, dorsal view; **B.** Terminal segments of *Thrips menyanthidis*, female, lateral view; **C.** Habitus of *Chirothrips hamatus*, male, dorsal view (except for abdominal segments III–VII, which are shown in ventral view). Roman numbers indicate abdominal and antennal segment numbers.

Notes. Globally, the family Thripidae encompasses more than 2,000 described species (Mound et al. 2018), many known as significant agricultural pests, capable of transmitting plant viruses and causing direct feeding damage. Sweden hosts a subset of this diverse family, with numerous species adapted to the colder, northern climate. Thripidae species in Sweden exhibit diverse ecological niches, ranging from specialists found in floral habitats to more generalised species inhabiting agricultural and forest environments.

Key to the subfamilies and species of Thripidae from Sweden

1 Metathoracic endoskeletal furca lyre-shaped, extending to mesothoracic furca (Fig. 3A) ..... **Dendrothripinae**

– Metathoracic furca small, U- or Y-shaped, rarely extending to mesothoracic furca and then only as a slender spinula (Fig. 2A) **2**

2(1) Head with conspicuous honeycomb-patterned reticulation; always macropterous; antennae with long, whiplike last segment; first vein of forewing more or less fused to costa (Fig. 4A)..... **Panchaetothripinae**

– Head not conspicuously strongly reticulated; doubtful species are apterous or with first vein and costa clearly separate..... **3**

3(2) Abdominal tergites covered with microtrichia, with at most a small bare patch medially ..... **Sericothripinae**

– Abdominal tergites without numerous microtrichia..... **Thripinae**

Dendrothripinae

1 Wing apex acute with a long terminal seta; antennae 7-segmented; minute yellow thrips with conspicuously dark wings (greenhouses/indoors) ..... ***Leucothrips nigripennis***

– Anterior margin of forewings recurved at apex to join posterior margin, terminal setae minute; antennae 8- or 9-segmented (Fig. 3B)..... **2**

2(1) Pronotum with one pair of long posteroangular setae..... ..... ***Dendrothrips saltator***

– Pronotum without prominent posteroangular setae **3**

3(2) Forewings uniformly shaded grey-brown (Fig. 3B)... ***Dendrothrips degeeri***

– Forewings banded (Fig. 3C) ..... ***Dendrothrips ornatus***

Panchaetothripinae

1 First vein of forewing with a complete row of stout setae (Fig. 4B); sense cones on antennal segment III and IV forked (Fig. 4C) (greenhouses/indoors)..... ***Hercinothrips femoralis***

– Forewing with only a few setae on first vein (Fig. 4D); sense cones on antennal segments III and IV simple **2**

2(1) Pronotum with one pair of long postero-angular setae; forewings broad with wing membrane reticulate (Fig. 4A) (greenhouses/indoors)..... ..... ***Parthenothrips dracaenae***

– Pronotum without prominent postero-angular setae; forewings slender without reticulate pattern (Fig. 4D) (greenhouses/indoors)..... ..... ***Heliothrips haemorrhoidalis***

Figure 3. Dendrothripinae. **A.** *Dendrothrips degeeri*, mesothoracic furca; **B.** *Dendrothrips degeeri*, female, habitus; **C.** *Dendrothrips ornatus*, female, in vivo.

Sericothripinae

1 Abdominal tergites medially with a bare patch lacking microtrichia (Fig. 4E); metanotum without microtrichia; macropterous ..... ***Neohydatothrips gracilicornis***

– Abdominal tergites completely covered with microtrichia; metanotum with microtrichia near posterior margin at least; micropterous or macropterous **2**

2(1) Microtrichia on metanotum forming a transverse band at posterior third (Fig. 4F) ..... ***Sericothrips bicornis***

– Metanotal microtrichia irregularly arranged, often forming a triangular patch covering posterior half (Fig. 4G) ..... ***Sericothrips staphylinus***



Figure 4. A–D. Panchaetothripinae; **A.** *Parthenothrips dracaenae*, female, head, pronotum and wing; **B.** *Hercinothrips femoralis*, wings; **C.** *H. femoralis*, antenna; **D.** *Heliothrips haemorrhoidalis*, wings; **E–G.** Sericothripinae; **E.** *Neohydatothrips gracilicornis*, abdominal tergites; **F.** *Sericothrips bicornis*, meso- and metanotum; **G.** *S. staphylinus*, meso- and metanotum.

Thripinae

1 Abdominal tergites with microtrichia, at least some microtrichia present laterally along the lines of sculpture or marginally (Fig. 5A) and tergal ctenidia always lacking (other comblike structures might be present) **2**

– Abdominal tergites without microtrichia; ctenidia present (Fig. 2A, B) or absent **7**

2(1) Sternites with microtrichia **3**

– Sternites without microtrichia **4**

3(2) Antennae 8-segmented (greenhouses/indoors) ..... ***Scirtothrips longipennis***

– Antennae 6-segmented ..... ***Drepanothrips reuteri***

4(2) Forewing with 2 visible longitudinal veins (second vein bearing at least 2 setae) (Fig. 5B); maxillary palps 3-segmented (Fig. 5B, C) **5**

– Forewing apparently with only one longitudinal vein (second vein without setae); maxillary palps 2-segmented (Fig. 9C) (greenhouses/indoors) ..... ***Echinothrips americanus***

5(4) Ocellar setae s3 situated medially in front of hind ocelli (Fig. 5E); mesothoracic furca weak or absent (Fig. 5G) ..... ***Mycterothrips latus***

– Ocellar s3 situated between hind ocelli; both meso- and metathoracic furca with spinula (Fig. 5B) **6**

6(5) Postocular setae s1 conspicuously longer than s3 (Fig. 5F) ..... ***Mycterothrips consociatus***

– Postocular setae s3 and s1 subequal in length (Fig. 5D) ..... ***Mycterothrips salicis***

7(1) Metathoracic furca with spinula (Fig. 5B); head, pronotum, and wings bearing conspicuously long setae (Fig. 5H) ..... ***Scolothrips uzeli***

– Metathoracic furca without spinula (Fig. 2A) **8**

8(7) Ctenidia present, at least on abdominal tergites VII–VIII (Fig. 2A, B) **9**

– Abdominal tergites without ctenidia, but sometimes with a group of irregular microtrichia **49**

9(8) Maxillary palps 3-segmented (Fig. 5C) **10**

– Maxillary palps 2-segmented (Fig. 9C) **47**

10(9) Antennal segments III and IV with forked sense cones; mesothoracic spinula present (Fig. 2A) **11**

– Antennal segments III and IV with simple sense cones; mesothoracic spinula absent (Fig. 5G) **48**

11(10) Ctenidia on tergite VIII laterad in front of spiracles (Fig. 5I); head with 2 pairs of anteocellar setae (Fig. 5J, N) **12**

– Ctenidia on tergite VIII mesad and posterior to spiracles (Fig. 6A); head with a single pair of anteocellar setae (s1 absent) (Fig. 6B) **17**

12(11) Fore tarsi at least with a small tooth (Fig. 5K, L); interocellar setae usually arising between hind ocelli (Fig. 5J) **13**

– Fore tarsi unarmed; interocellar setae usually arising in front of hind ocelli **14**

13(12) Fore tarsus apically with a curved apical claw (Fig. 5K); antennal segment V rounded..... ***Kakothrips priesnerorum***

– Fore tarsus apically with a triangular tooth (Fig. 5L); antennal segment V slender..... ***Kakothrips pisivorus***

14(11) Metanotum without campaniform sensilla (Fig. 5M)..... **15**

– Metanotum with campaniform sensilla (Fig. 5P) **16**

15(14) Head projecting in front of eyes; cheeks straight and approx. as long as compound eyes (Fig. 5N) ..... ***Frankliniella tenuicornis***

– Head not projecting; cheeks rounded and shorter than compound eyes (Fig. 5O) ..... ***Frankliniella intonsa***

16(14) All postocular setae s4 not very long and stout (Fig. 5Q); pronotum usually with 2 (occasionally 3) anteromarginal setae ..... ***Frankliniella pallida***

– Postocular setae s4 long and stout (Fig. 5R); pronotum usually with 4 (3–5) anteromarginal setae (greenhouses/indoors)..... ***Frankliniella occidentalis***

17(11) Tibiae with 3 long setae (Fig. 6C); abdominal sternite II with 6 marginal setae (middle pair slightly anterior to margin) (Fig. 6D); ferna separated (Fig. 6E) ..... ***Baliothrips kroli***

– Tibiae without distinctive long setae (Fig. 6F); abdominal sternite II with 4 marginal setae (Fig. 6G); ferna fused (Fig. 6H) **18**

18(17) At least abdominal sternites III–IV with accessory setae in addition to the 6 marginal setae (Fig. 6G)..... **19**

– All abdominal sternites without accessory setae..... **30**

19(18) Pleurotergites III–IV with accessory setae medially (Fig. 6I) **20**

– Pleurotergites III–IV with setae at posterior margin only..... **24**

20(19) Antennae 8-segmented (Fig. 6J) **21**

– Antennae 7-segmented (Fig. 7A) **23**

21(20) Forewing pale; first vein with 3 setae on distal half of wing (Fig. 6K); antennal segment III yellow (Fig. 6J)..... ***Thrips vulgatissimus***

– Forewings shaded; first vein with at least 5 setae on distal half of wing (Fig. 6M); antennal segment III at least shaded brown (Fig. 6L) **22**

22(21) Antennal segment III slender; as dark as IV (Fig. 6L); tergite VIII in females with marginal comb of microtrichia complete (Fig. 6N) ..... ***Thrips atratus***

– Antennal segment III swollen; lighter than IV; tergite VIII in females with marginal comb of microtrichia interrupted medially (Fig. 7E) ..... ***Thrips linariae***

23(20) Forewing with 3 setae on distal half of first vein (Fig. 6K); abdominal tergite II with 3 lateral setae (Fig. 7B); accessory setae arranged in 2 rows, at least on posterior sternites (Fig. 7C) ..... ***Thrips pillichi***

– Forewing with 7–11 setae on distal half of first vein (Fig. 6M); abdominal tergite II with 4 lateral setae (Fig. 7F); accessory setae arranged in 1 row (Fig. 7D)..... ***Thrips minutissimus***

24(19) Antennae 8-segmented (Fig. 6J) **25**

– Antennae 7-segmented (Fig. 7A) **26**

25(24) Fore wing with 3 setae on distal half of first vein (Fig. 6K); tergite II with 4 setae on lateral margin (Fig. 7F) ..... ***Thrips pini***

– Fore wing with more than 5 setae on distal half of first vein; tergite II with 3 setae on lateral margin (Fig. 7B) (greenhouses/indoors) ..... ***Thrips simplex***

26(24) Accessory setae arranged in more than 1 transverse row (Fig. 7C); frequently micropterous or hemimacropterous; macropterous forms with 7–11 setae on distal half of first wing vein ..... ***Thrips angusticeps***

– Accessory setae arranged in a single transverse row (Fig. 7D); macropterous, forewing with 3 or 4 setae on distal half of first vein **27**

27(26) Sternite III – VI with 1–5 accessory setae each; pleurotergites with rows of microtrichia (Fig. 7G); female tergite IX with 1 pair of campaniform sensilla (anterior pair absent) (Fig. 7H) ..... ***Thrips origani***

– Sternal accessory setae more numerous; pleurotergites without ciliate microtrichia; tergite IX with 2 pairs of campaniform sensilla (Fig. 7L) **28**

28(27) Fore tarsus with a slender claw at apex (Fig. 7I); metanotum with elongate reticulations medially (Fig. 7J) ..... ***Thrips calcaratus***

– Fore tarsus unarmed; metanotum more or less striate medially (Fig. 7K) **29**

29(28) Male brown: female x + 2y + z > 430 µm (where x = length of inner pronotal posteroangular setae (Fig. 7M), y = length of median setae (s1) on abdominal tergite IX, z = length of abdominal tergite X (Fig. 7L)) .... ***Thrips trehernei***

– Male yellow; female with x + 2y + z < 430 µm (see notes above) ..... ***Thrips physapus***

30(18) Pleurotergites III – IV with one or more accessory setae medially (Fig. 6I) **31**

– Pleurotergites III – IV with setae at posterior margin only **32**

31(30) Body colour yellow; setae dark; metanotum very closely striate ..... ***Thrips brevicornis***

– Body colour brown; setae dark; legs and antennal segment III yellow; metanotum closely striate ..... ***Thrips fulvipes***

32(30) Abdominal tergite VIII with complete comb of microtrichia on posterior margin (Fig. 6N) **33**

– Either abdominal tergite VIII without posterior marginal comb, or comb interrupted medially (Fig. 7E) or reduced to broad triangular teeth (Fig. 7T) **40**

33(32) Abdominal pleurotergites with rows of microtrichia (Fig. 7G); tergite IX in females with 1 pair of campaniform sensilla (anterior pair absent) (Fig. 7H); males with glandular areas on sternites III – V; forewing usually with 4 (sometimes 3–5) setae on distal half of first vein ..... ***Thrips tabaci***

– Abdominal pleurotergites without rows of microtrichia; tergite IX in females frequently with 2 pairs of campaniform sensilla (Fig. 7L); males with glandular areas on sternites III–VII (Fig. 7N); forewing usually with 3 setae on distal half of first vein (Fig. 6K) **34**

34(33) Head and thorax yellow, or if yellow with faint brown markings, then median pairs of tergal setae (s1, s2) half as long as tergites length (Fig. 7Q) **35**

– Head and thorax brown **39**

35(34) Interocellar setae arising posterior to fore ocellus, within ocellar triangle (Fig. 7O); body colour yellow without markings; antennal segments IV – V bicoloured ..... ***Thrips flavus***

– Interocellar setae arising almost laterad of fore ocellus, anterolateral of ocellar triangle (Fig. 7P) **36**

36(35) Tergites IV – V with 2 pairs of median setae (s1, s2) half as long as tergites length (Fig. 7Q); body colour yellow with faint brown markings; macropterous, hemimacropterous or macropterous ..... ***Thrips nigropilosus***

– Median setae on tergites IV – V < 0.3 × as long as length of these tergites; head and thorax yellow, abdominal tergites sometimes with grey markings medially **37**

37(36) Abdominal tergite IX of female with 1 pair of campaniform sensilla (anterior pair absent) (Fig. 7H); metanotal sculpture reticulate; abdominal tergites yellow, usually with grey markings medially ..... ***Thrips urticae***

– Abdominal tergite IX of females with 2 pairs of campaniform sensilla (Fig. 7L); metanotal sculpturing striate; abdominal tergites yellow without markings medially **38**

38(37) Setae s2 on tergites II – V light; tergite V with setae s2 clearly shorter and weaker than s3 (Fig. 7R) ..... ***Thrips alni***

– Setae s2 on tergites II – V dark brown; tergite V with setae s2 and s3 subequal in length and strength (greenhouses/indoors) ..... ***Thrips palmi***

39(34) Tergite II with 3 setae on lateral margin (Fig. 7B); antennal segments III and IV swollen with constricted vasiform apices (Fig. 7S); females rarely macropterous, usually micropterous or hemimacropterous; males micropterous, with transverse slender pore plates on sternites III – VII ..... ***Thrips dilatatus***

– Tergite II with 4 setae on lateral margin (Fig. 7F); antennal segments III and IV without an elongated apical neck; males with small circular pore plates on sternites III–VII ..... ***Thrips validus***

40(32) Females usually micropterous, rarely macropterous; metanotum medially reticulate; tergite VIII with comb on posterior margin represented by a few weak triangular teeth only (Fig. 7T); males micropterous;. **41**

– Both sexes macropterous; metanotum with longitudinal sculpture medially **42**

41(40) Antennal segments III and IV swollen with constricted vasiform apices (Fig. 7S); tergite II with 3 setae on lateral margin (Fig. 7B) ..... **see above: *Thrips dilatatus* 39**

– Antennal segments III and IV without an elongated apical neck; tergite II with 4 setae on lateral margin (Fig. 7F); females with head and thorax yellow or light-brown in contrast to dark brown abdomen; males yellow, with transverse oval pore plates on sternites III–VII ..... ***Thrips discolor***

42(40) Pronotum with lines of sculpture absent or very faint; males with pore plates on sternite III – VI ..... ***Thrips viminalis***
– Pronotum with many well-developed transverse lines of sculpture (Fig. 7U); males with pore plates on sternite III–VII (Fig. 7N).....**43**
43(42) Tergite II with 3 setae on lateral margin (Fig. 7B); lines of sculpture on pleurotergites with ciliate microtrichia (Fig. 7G).....**44**
– Tergite II with 4 setae on lateral margin (Fig. 7F); lines of sculpture on pleurotergites with dentate microtrichia (Fig. 7V).....**45**
44(43) Abdominal sternite I with 2 or 3 microsetae on the anterior margin (Fig. 7X); pronotal posteroangular setae short (usually < 46 µm)..... .....***Thrips juniperinus***
– Abdominal sternite I without microsetae on anterior margin; pronotal postero-angular setae long (usually > 54 µm) .....***Thrips major***
45(43) Fore wings shaded, pale only at base (Fig. 7W) **46**
– Fore wings clear, uniformly pale throughout their length (Fig. 8A)..... .....***Thrips roepkei***
46(45) Median postocular setae (s1) slightly closer together than posterior ocelli; antennal segment III as dark as distal segments (Fig. 8B); males brown.....***Thrips menyanthidis***
– Median postocular setae (s1) approximately as far apart as posterior ocelli; antennal segments IV and V darker than III (Fig. 8C); males yellow.....***Thrips fuscipennis***
47(9) Mesothoracic furca with spinula (Fig. 2A); abdominal sternite II with 4 setae .....***Stenothrips graminum***
– Mesothoracic furca without spinula; abdominal sternite II with 6 setae ..... ***Baliothrips dispar***
48(10) Antennae 8-segmented; body brown; sternites without discal setae.... ..... ***Iridothrips iridis***
– Antennae 7-segmented; body yellow with only segment X brown (Fig. 8D); sternal discal setae present..... ***Bolacothrips jordani***
49(8) Pronotum with 2 pairs of postero-angular setae (Fig. 2A).....**50**
– Pronotum with 1 pair or without postero-angular setae **68**
50(49) Head usually with 2 pairs of ante-ocellar setae (Fig. 8E).....**51**
– Head with only 1 pair of ante-ocellar setae or ocellar setae lacking.....**61**
51(50) External sense cone on antennal segment VI with elongated base (Fig. 8H); fore tibia with distal teeth (Fig. 8F).....**52**
– External sense cone on antennal segment VI round at base; fore tibia without distal teeth **57**
52(51) Fore tibia with short teeth, not longer than the setae (Fig. 8F)..... .....***Odontothrips confusus***
– Fore tibia with 1 or 2 large apical claws (Fig. 8G, 8J).....**53**
53(52) Fore tibiae with 1 apical claw (Fig. 8G) **54**
– Fore tibiae with 2 apical claws (Fig. 8J).....**56**
54(53) Extruding part of sense cone of antennal segment VI short, at most 6 µm (Fig. 8H); upper vein in forewing with a short interval of setae missing in a row (like 1 seta missing).....***Odontothrips loti***
– Extruding part of sense cone of antennal segment VI long, at least 10 µm (Fig. 8I); upper vein in forewing with a long interval of setae missing in a row (like 4–7 setae missing) **55**

55(54) Margins of antennal segment III basoventrally expanded (Fig. 8I) ..... ***Odontothrips phaleratus***

– Margins of antennal segment III evenly symmetrical..... ***Odontothrips intermedius***

56(53) Fore tarsi with 1 or 2 small protuberances (Fig. 8J)..... ***Odontothrips biuncus***

– Fore tarsi smooth..... ***Odontothrips ulicis***

57(51) Posterior margin of abdominal tergites II–VII with craspeda or at least with rounded or triangular plates (Fig. 8K, M)..... **58**

– Posterior margin of abdominal tergites without craspeda or other projections; sternite VII in females with median setae (s1) arising just anterior to posterior margin; males apterous, with three pore plates on each of sternites III–VIII ..... ***Ceratothrips ericae***

58(57) Pronotum rectangular; mesothoracic furca with spinula (Fig. 8L) ..... ***Firmothrips firmus***

– Pronotum trapezoidal; mesothoracic furca without spinula (Fig. 2C)... **59**

59(58) Antennal segment II evenly symmetrical ..... ***Chirothrips hamatus***

– Antennal segment II produced outwards or at least asymmetrical (Fig. 8N)..... **60**

60(59) Tergite VII with campaniform sensilla posterior to subbasal line (Fig. 8M) ..... ***Chirothrips manicatus***

– Tergite VII with campaniform sensilla anterior to subbasal line (Fig. 8O) ..... ***Chirothrips ambulans***

61(50) Antennae 7-segmented; abdominal tergites with setae s1–s3 strong . ..... ***Platythrips tunicatus***

– Antennae 8-segmented **62**

62(61) Fringes at anterior margin of forewing always longer than costal setae; postocular setae s2 short (Fig. 8P); antennal segment IV without an additional sense cone **63**

– Costal setae of forewing longer than fringes; postocular setae s2 conspicuously long; “antennal segment III five times longer than wide, antennal segment IV with an additional simple sense cone (greenhouses/indoors)..... ***Bradinothrips musae***

63(62) Antennal segments VII–VIII together longer than VI; segment VI broadly attached to V (Fig. 8P) ..... ***Rhaphidothrips longistylosus***

– Terminal antennal segments shorter; segment VI not broadly attached to V..... **64**

64(63) Pterothorax and abdomen with polygonal reticulate sculpture (Fig. 8Q); postocular setae distinctly longer than interocellar setae ..... ***Ctenothrips distinctus***

– Pterothorax and abdomen without polygonal sculpture; postocular setae shorter than interocellar setae..... **65**

65(64) Body dark; other characters differing from below **66**

– Body yellow; forewings pale with brown cross-bands (Fig. 8R); abdominal tergite VIII with stippled area of sculpture extending medially from spiracles (greenhouses/indoors) ..... ***Chaetanaphothrips orchidii***

66(65) Hind margin of abdominal tergites without craspeda; tergite VIII with marginal comb present (Fig. 6N); macropterous.....**67**

– Hind margin of abdominal tergites with short craspeda; tergite VIII without marginal comb; micropterous (macropterous forms not known from Sweden) (Fig. 9A) ..... ***Tmetothrips subapterus***

67(66) Fore tarsi with terminal claw (Fig. 9B); first vein with 3–8 distal setae ..... ***Taeniothrips inconsequens***

– Fore tarsi without terminal claw; first vein with 1 + 2 distal setae..... ..... ***Taeniothrips picipes***

68(49) Hind margin of pronotum with 4 pairs of posteromarginal setae **69**

– Hind margin of pronotum with 2 pairs of posteromarginal setae; posteroangular setae stout. long (up to 75 µm) and spatulate .... ***Pteridothrips pteridicola***

69(68) Pronotum with a pair of long posteroangular setae (Fig. 2A).....**70**

– Pronotum without conspicuous posteroangular setae.....**76**

70(69) Mesothoracic furca without spinula (Fig. 5G); maxillary palps 2-segmented (Fig. 9C); setae s1 on tergite X in females thorn-like (Fig. 9G) **71**

– Mesothoracic furca with spinula (Fig, 2A); maxillary palps 3-segmented (Fig. 5C); setae s1 on tergite X in females normal, bristle-like.....**74**

71(70) Antennal segment II or III strongly asymmetric (Fig. 9D).....**72**

– Antennal segment II or III not strongly prolonged externally (Fig. 9E) **73**

72(71) Antennal segment III prolonged externally (Fig. 9D) .... ***Limothrips denticornis***

– Antennal segment II prolonged apically..... ***Limothrips angulicornis***

73(71) Antennal segments III–IV slightly asymmetric (Fig. 9E); tarsi 1-segmented (Fig. 9F); abdominal tergite IX in females without thorn-like setae (Fig. 9G)..... ***Limothrips cerealium***

– Antennal segments III–IV symmetric (Fig. 9H); tarsi 2-segmented (Fig. 9I); abdominal tergite IX in females with strong thorn-like setae (Fig. 9J)..... ***Limothrips consimilis***

74(70) Fore tarsi with a curved apical claw (Fig. 9K) ..... ***Oxythrips ajugae***

– Fore tarsi without protrusions.....**75**

75(74) Abdominal sternites III–VI with discal setae (Fig. 9L) .... ***Oxythrips bicolor***

– Abdominal sternites III–VI without discal setae..... ..... ***Oxythrips ulmifoliorum***

76(69) Antennal style (segments VII–VIII) approx. as long as or slightly longer than segment VI **77**

– Antennal style (segments VII–VIII) much shorter than segment VI, or style absent (antennae thus 6-segmented) **78**

77(76) Antennal segment III paler than remaining segments, as pale as fore tibia; abdominal segment X in females tapering posteriorly to a conical apex (Fig. 9M) ..... ***Belothrips morio***

– Antennal segment III brown, darker than fore tibia (Fig. 9N); side margins in distal half of abdominal segment X in females subparallel..... ..... ***Belothrips acuminatus***

78(76) Abdominal tergites with craspeda (Fig. 8K) **79**
– Marginal craspeda on abdominal tergites not developed **81**
79(78) Mesothoracic furca without spinula; body without reticulate sculpture; apterous **80**
– Mesothoracic furca with spinula (Fig. 2A); head, thorax, and legs with reticulate sculpture; macropterous ..... ***Dictyothrips betae***
80(79) Body slender, yellow; males (Fig. 9O) ..... **males of *Aptinothrips* 82**
– Body stout; dark ..... ***Apterothrips secticornis***
81(78) Head longer than wide; antennal segments III–IV with simple sense cones (Fig. 9P); mesothoracic furca without spinula; apterous **82**
– Head wider than long; antennal segments III–IV with forked sense cones (Fig. 10A); mesothoracic furca with spinula; macropterous or macropterous **84**
82(81) Antennae 6-segmented; abdominal sternites usually with discal setae (Fig. 9Q) **83**
– Antennae 8-segmented; abdominal sternites without discal setae ..... ***Aptinothrips stylifer***
83(82) Outline of antennal segment II oval, widest at mid-length; petiole exposed (Fig. 9P); posteromarginal setae on sternites III–IV arising at margin; sternites of males without pore plates ..... ***Aptinothrips rufus***
– Outline of antennal segment II dome-shaped, widest at base; petiole concealed; posteromarginal setae on sternites III–IV arising in front of margin; sternites III–VI of males with small circular pore plates ..... ***Aptinothrips elegans***
84(81) Median setae (s1) at least on tergites VI–VIII long, longer than their distance apart (Fig. 10B) **85**
– Median setae (s1) on tergites VI–VIII short and weak, shorter than their distance apart **88**
85(84) Median setae on tergites III–VI close to subbasal line (Fig. 10B); males with pore plates on sternites III–VI **86**
– Median setae (s1) on tergites III–VI at least 20 μm posterior of subbasal line (Fig. 10C); males with pore plates on sternites III–VII ..... ***Belothrips ferrugineus***
86(85) Body yellow; median brown mottling occupying approx. half of the tergites; antennal segment I yellow; posterior margin of head brown ..... ***Belothrips sordidus***
– Body, head, and antennal segment I brown **87**
87(86) Metanotum with campaniform sensilla (Fig. 10D); thoracic dorsal setae rod-like ..... ***Belothrips validus***
– Metanotum without campaniform sensilla; thoracic dorsal setae finely pointed ..... ***Belothrips silvarum***
88(84) Clavus of forewing with 5 venal setae (Fig. 10E, F) or macropterous **89**
– Macropterous; clavus of forewing with 6 venal setae (greenhouses/indoors) ..... ***Aurantothrips orchidaceus***
89(88) Body dark brown; antennal segment I as dark as II (Fig. 10F); macropterous ..... ***Anaphothrips badius***
– Body yellow or yellowish brown; antennal segment I lighter than II; macropterous or macropterous ..... ***Anaphothrips obscurus***

Checklist

This checklist comprehensively lists all known Swedish Thripidae species (including stable and reproducing or more limited and sporadic), with new national and regional records, providing details on their provincial distribution. The examined material is presented for new records, both national and provincial.



Figure 5. **A–D.** *Mycterothrips salicis*; **A.** Abdominal tergites with microtrichia on lateral sculpture lines; **B.** Forewing, head and thorax (ventral side); **C.** Mouthparts; **D–F.** Head; **D.** *M. salicis*; **E.** *M. latus*; **F.** *M. consociatus*; **G.** *M. latus*, meso- and metathoracic furca; **H.** *Scolothrips uzeli*, habitus; **I–K.** *Kakothrips priesnerorum*; **I.** Abdominal tergite VIII; **J.** Head, s1 anteocellar seta 1; **K.** Foretarsus; **L.** *K. pisivorus*, foretarus; **M, N.** *Frankliniella tenuicornis*; **M.** Metanotum; **N.** Head; **O.** *F. intonsa*, head; **P, Q.** *F. pallida*; **P.** Metanotum; **Q.** Head. R. *F. occidentalis*, head, s4 postocular seta 4. Abbreviations: sc = campaniform sensilla.

Subfamily Dendrothripinae Priesner, 1925

Dendrothrips degeeri Uzel, 1895

Distribution. Palearctic. Nordic distribution: Sweden, Norway, Finland. Swedish distribution: Sk, Öl, Sm, Go, Ds, Sö, Up.

Remarks. Feeding and breeding on leaves of various deciduous trees and herbaceous plants.

Dendrothrips ornatus (Jablonowski, 1894)

Distribution. Palearctic, introduced in Nearctic. Nordic distribution: Sweden, Norway, Finland. Swedish distribution: Sö.

Remarks. First record for Sö, previous records not specified. Living on the leaves of *Fraxinus*, *Ligustrum*, *Syringa*, as well as *Alnus*, *Corylus*, and *Tilia*, here collected on *Tilia cordata* (Malvaceae).

Material examined. SWEDEN • 5♀♀; Sö, Nyköpings kommun, Tovetorp research station, 58.94788°N, 17.14470°E, on *Tilia cordata*; 2024.viii.28; E. Wahlberg leg.; voucher IDs: DA2T, DA8T, DA9T, DB1T, DE7T.

Dendrothrips saltator Uzel, 1895

Distribution. Palearctic. Nordic distribution: Sweden, Norway, Finland. Swedish distribution: Sk, Ds, Vr.

Remarks. First record for Sk. Feeding and breeding on a wide range of herbaceous plants and trees, both coniferous and deciduous.

Material examined. SWEDEN • 1♀; Sk, Lunds kommun, Botaniska trädgården, 55.705833°N, 13.204167°E, on *Origanum vulgare*, 2025.viii.21, leg. C.-A. Gertsson.

Leucothrips nigripennis Reuter, 1904

Distribution. Nearctic, but introduced worldwide, in temperate regions indoors and in greenhouses. Nordic distribution: Sweden, Denmark, and Finland. Swedish distribution: sporadically indoors/in greenhouses.

Remarks. Feeding and breeding on different species of ferns.

Subfamily Panchaetothripinae Bagnall, 1912

Heliothrips haemorrhoidalis (Bouché, 1833)

Distribution. Neotropical, but introduced worldwide in tropical areas, in temperate areas in greenhouses and indoors. Nordic distribution: Sweden, Norway, Denmark, Finland. Swedish distribution: Sk, Sm, Sö. Up.

Remarks. Feeding and breeding on both coniferous and deciduous trees and herbaceous plants, often in water-stressed environments.



Figure 6. A–E. *Baliothrips kroli*; **A.** Tergite VIII; **B.** Head; **C.** Midtibia and -tarsus; **D.** Abdominal sternites I–II; **E.** Prosternum; **F–K.** *Thrips vulgatissimus*; **F.** Midtibia- and tarsus; **G.** Abdominal sternites II and III; **H.** Prosternum; **I.** Pleurotergite; **J.** Antenna; **K.** Forewing; **L–N.** *T. atratus*; **L.** Antenna; **M.** Forewing; **N.** Female abdominal tergite VIII.

***Hercinothrips femoralis* (O. M. Reuter, 1891)**

Distribution. Afrotropical, but introduced worldwide in tropical and subtropical areas, in temperate areas in greenhouses and indoors. Nordic distribution: Sweden, Norway, Denmark, Finland. Swedish distribution: Sk, Up, Vb.

Remarks. Breeding and feeding on leaves of different herbaceous plants.



Figure 7. **A–C.** *Thrips pillichi*; **A.** Antenna; **B.** Abdominal tergite II; **C.** Abdominal sternite VII; **D.** *T. minutissimus*, abdominal sternite VII; **E.** *T. linariae*, abdominal tergite VIII; **F.** *T. pini*, abdominal tergite II; **G, H.** *T. origani*; **G.** Pleurotergite; **H.** Female abdominal tergites VIII–X; **I, J.** *T. calcaratus*; **I.** Fore tarsus; **J.** Metanotum; **K–M.** *T. trehernei*; **K.** Metanotum; **L.** Female abdominal tergites IX–X; **M.** Pronotum; **N, O.** *T. flavus*; **N.** Male abdominal sternites III–VIII; **O.** Head; **P, Q.** *T. nigropilosus*; **P.** Head; **Q.** Abdominal tergite II; **R.** *T. alni*, abdominal tergite V; **S, T.** *T. dilatatus*; **S.** Antenna; **T.** Female abdominal tergites VIII–IX; **U.** *T. major*, pronotum; **V, W.** *T. menyanthidis*; **V.** Pleurotergites; **W.** Forewing; **X.** *T. juniperinus*, abdominal sternite I. Abbreviations: cs = campaniform sensilla, ios = interocellar seta, pa = posteroangular seta, s1 = median seta, s2 = setae 2, s3 = setae 3.

Parthenothrips dracaenae (Heeger, 1854)

Distribution. Afrotropical, but introduced worldwide in tropical and sub-tropical regions, in temperate areas mainly in greenhouses, but has been collected outdoors in Norway (Kobro and Ultizka 2021). Nordic distribution: Sweden, Norway, Denmark, Finland, Iceland. Swedish distribution: Sk, Sm, Bo, Up, Vr.

Remarks. Feeding and breeding on, in majority, hard-leaved plants.

Figure 8. **A.** *T. roepkei*, forewing; **B.** *T. menyanthidis*, head and partial antenna; **C.** *T. fuscipennis*, head and partial antenna; **D.** *Bolacothrips jordani*, apterous female, habitus; **E, F.** *Odontothrips confusus*; **E.** Head; **F.** Foretibia and -tarsus; **G, H.** *O. loti*; **G.** Foretibia and -tarsus; **H.** Apical antennal segments; **I.** *O. phaleratus*, antenna; **J.** *O. biuncus*, foretibia and -tarsus; **K, L.** *Firmothrips firmus*; **K.** Abdominal tergite VII; **L.** Micropterous female, habitus; **M, N.** *Chirothrips manicatus*; **M.** Abdominal tergite VII; **N.** Antenna; **O.** *C. ambulans*, abdominal tergite VII; **P.** *Rhaphidothrips longistylosus*, head and antennae; **Q.** *Ctenothrips distinctus*, meso- and metanotum; **R.** *Chaetanaphothrips orchidii*, forewing. Abbreviations: as = anterocellar setae, cs = campaniform sensilla, ps1 = postocular seta 1, sc = sense cone on segment VI.

Subfamily Sericothripinae Karny, 1921

***Neohydatothrips gracilicornis* (Williams, 1916)**

Distribution. Palearctic. Nordic distribution: Sweden, Norway, Finland. Swedish distribution: Sk, Öl, Go, Ög, Vg, Vr, Sö, Up, Gä, Me, Hr, Vb, To.



Figure 9. **A.** *Tmetothrips subapterus*, female, habitus; **B.** *Taeniothrips inconsequens*, foretarsus; **C, D.** *Limothrips denticornis*; **C.** Mouthparts; **D.** Antenna; **E–G.** *L. cerealium*; **E.** Antenna; **F.** Foretarsus; **G.** Female abdominal tergites VIII–X; **H–J.** *L. consimilis*; **H.** Antenna; **I.** Foretarsus; **J.** Female abdominal tergite IX; **K.** *Oxythrips ajugae*, foretarsus; **L.** *O. bicolor*, sternite V; **M.** *Belothrips morio*, female habitus; **N.** *B. acuminatus*, antenna; **O–Q.** *Aptinothrips rufus*; **O.** Male habitus; **P.** Antenna; **Q.** Abdominal sternite V.

Remarks. First records for Up, Gä, and Me. Preferably in warmer places. This species is mainly found on *Vicia* sp., particularly on *Vicia cracca*; occasionally also on leaves of deciduous trees.

Material examined. SWEDEN • 1♂; Up, Uppsala kommun, Lågbol, 60.034883°N, 18.420316°E, on *Calamagrostis*; 2021.vii.19; E. Wahlberg leg.; voucher ID: AP4T. • 2♀♀; Gä, Gävle kommun, Östra Sikvik, 60.66559°N, 17.29876°E, mixed herbaceous vegetation; 2023.vi.20; E. Wahlberg leg.; voucher IDs: DL1T, DL2T. • 2♀♀; Me, Timrå kommun, Indalsälvens delta, 62.508667°N, 17.448164°E, mixed herbaceous vegetation; 2023.vi.14; E. Wahlberg leg.; voucher IDs: DJ5T, DJ6T.



Figure 10. A, B. *Belothrips validus*. Antennal segment IV; **B.** Abdominal tergite IV; **C.** *B. ferrugineus*, abdominal tergite IV; **D.** *B. validus*, metanotum; **E, F.** *A. badius*; **E.** Forewing clavus; **F.** Female habitus. Abbreviations: cs = campaniform sensilla, s1 = seta 1, sc = sense cone.

Sericothrips bicornis (Karny, 1910)

Distribution. Palearctic. Nordic distribution: Sweden, Denmark, Finland. Swedish distribution: Sk, Öl, Sö, Up, Ly.

Remarks. Feeding and breeding on leaves of Fabaceae, mainly on creeping species like *Lotus corniculatus* and *Trifolium repens*.

Sericothrips staphylinus Haliday, 1836

Distribution. Palearctic, introduced in other parts of the world as biological control agent. Nordic distribution: Sweden, Denmark. Swedish distribution: Sm, Öl, Ög, Bo, Sö, Up, Hälsningland.

Remarks. Feeding and breeding on leaves of *Ulex*.

Subfamily Thripinae Stevens, 1829

Anaphothrips badius (Williams, 1913)

Distribution. Palearctic. Nordic distribution: Sweden, Norway, Denmark. Swedish distribution: Sk, Sm, Go, Sö, Up, Vr.

Remarks. First records for Sk, Sm, and Sö. Feeding and breeding on leaves of Poaceae and Cyperaceae, here also collected on *Rubus* (Rosaceae). Prefers wet meadows and other damp areas.

Material examined. SWEDEN • 2♀♀; Sk., Kristianstads kommun, Gropahålet, Helge river mouth, 55.859063°N, 14.235729°E, on *Carex acuta*, 2024.v.05; E. Wahlberg leg.; voucher IDs: FF1T, FF2T • 1♀; Sk., Kristianstads kommun, Gropahålet, Helge river mouth, 55.859063°N, 14.235729°E, on *Leymus arenarius*, 2024.v.05; E. Wahlberg leg.; voucher ID: FE3T • 1♀; Sk., Kristianstads kommun, Lyngsjö nature reserve, 55.932709°N, 14.067986°E, wetland, 2024.v.21; E. Wahlberg leg.; voucher ID: FG6T • 2♀♀; Sm., Västerviks kommun, Hallmare, 57.870555°N, 16.756545°E, on *Phragmites australis*, 2024.v.05; E. Wahlberg leg.; voucher IDs: ES7T, ES8T. • 1♀; Sm., Västerviks kommun, Hallmare, beach, 57.863499°N, 16.760644°E, on *Bolboschoenus maritimus*, 2024.v.05; E. Wahlberg leg.; voucher ID: EA6T. • 1♀; SE, Sm., Västerviks kommun, Risebo nature reserve, pond edge, 58.031361°N, 16.106291°E, 2024.v.28; E. Wahlberg. leg; voucher ID: EZ9T. • 1♂; Sö, Nyköpings kommun, Tovetorp research station, 58.9476°N, 17.1467°E, on *Rubus idaeus*; 2024.viii.27; E. Nein leg.; voucher ID: DA7T.

Anaphothrips obscurus (O.F. Müller, 1776)

Distribution. Global in temperate regions. Nordic distribution: Sweden, Norway, Denmark, Finland. Swedish distribution: Sk, Bl, Sm, Go, Ög, Vg, Bo, Sö, Up, Vs, Vr, Dr, Me.

Remarks. First records for Bl and Me. Feeding and breeding on leaves of Poaceae.

Material examined. SWEDEN • 4♀♀; Bl, Karlskrona kommun, Knösö, 56.165360°N, 15.666374°E, on *Phragmites*; 2023.v.30; E. Wahlberg leg.; voucher IDs: CV2T, CV4T, CV5T, CV6T. • 4♀♀; Bl, Karlskrona kommun, Augerum, 56.217489°N, 15.673053°E, on Poaceae; 2023.v.30; E. Wahlberg leg.; voucher IDs: CX4T, CX5T, CX6T, CX8T. • 3♀♀; Me, Timrå kommun, Indalsälvens delta, 62.508667°N, 17.448164°E, in mixed herbaceous vegetation; 2023.vi.14; E. Wahlberg, leg.; voucher IDs: DJ4T, CC4T, CC5T.

Apterothrips secticornis (Trybom, 1896)

Distribution. Holarctic. Nordic distribution: Sweden, Norway, Finland, Iceland. Swedish distribution: Hr, Jä, Ly.

Remarks. Feeding and breeding on leaves of Poaceae and Cyperaceae.

Aptinothrips elegans Priesner, 1924

Distribution. Palearctic. Nordic distribution: Sweden, Norway, Denmark, Finland. Swedish distribution: Sö.

Remarks. Feeding and breeding on leaves of Poaceae.

Aptinothrips rufus Haliday, 1836

Distribution. Global in temperate regions. Nordic distribution: Sweden, Norway, Denmark, Finland, Iceland. Swedish distribution: Sk, Ha, Sm, Öl, Bo, Sö, Up, Dr, Nb.

Remarks. Feeding and breeding on leaves of Poaceae.

Aptinothrips stylifer Trybom, 1894

Distribution. Global. Nordic distribution: Sweden, Norway, Denmark, Finland, Iceland. Swedish distribution: Sk, Bl, Sm, Sö, Up, Vs, Vr, Hr, Jä, To.

Remarks. First record for Bl. Feeding and breeding on leaves of Poaceae.

Material examined. SWEDEN • 1♀; Bl, Karlskrona kommun, Knösö, 56.165360°N, 15.666374°E, on *Phragmites*; 2023.v.30; E. Wahlberg leg.; voucher ID: CV3T.

Aurantothrips orchidaceus (Bagnall, 1909)

Distribution. Neotropical, introduced in greenhouses worldwide on cultivated orchids. Nordic distribution: Sweden, Norway, Denmark (only indoors/in greenhouses). Swedish distribution: sporadically indoors/in greenhouses.

Remarks. Feeding and breeding on leaves of Orchidaceae.

Baliothrips dispar (Haliday, 1836)

Distribution. Holarctic. Nordic distribution: Sweden, Norway, Denmark, Finland. Swedish distribution: Vr, Dr, Jä, Vb, Ly.

Remarks. First records for Dr and Vb. Living in damp areas and feeding and breeding on leaves of Poaceae, Cyperaceae and *Typha* spp. (Typhaceae).

Material examined. SWEDEN • 2♀♀; Dr, Orsa kommun, Koppången nature reserve, wet land, 61.362499°N, 14.786772°E, on *Prunella vulgaris*, 2024.vii.30,; E. Wahlberg, leg.; voucher IDs: EF2T, EF3T. • 3♀♀; Vb, Skellefteå kommun, Nilsliden, 64.944032°N, 20.367689°E, mixed herbaceous vegetation; 2023.vi.18; E. Wahlberg, leg.; voucher IDs: CF5T, CG1T, CE8T.

Baliothrips kroli (Schille, 1911)

Distribution. Palearctic. Nordic distribution: Sweden, Norway, Denmark, Finland. Swedish distribution: Sk, Sm, Sö, Up.

Remarks. First record for Sö. Living in damp areas and feeding and breeding on leaves of Poaceae and Cyperaceae. Here also collected on *Sagittaria sagittifolia* (Alismataceae).

Material examined. SWEDEN • 2♀♀; Sö., Flens kommun, Sparreholm, beach, 59.086913°N, 16.829442°E, on *Carex acuta*, 2024.vii.29; E. Wahlberg, leg.; voucher IDs: EFE5, EFE6. • 1♀♀; Sö., Flens kommun, Sparreholms ekhagar, beach, 59.082452°N, 16.842549°E, on flowering *Sagittaria sagittifolia*, 2024.vii.25; E. Wahlberg, leg.; voucher IDs: EFJ7.

Belothrips acuminatus Haliday, 1836

Distribution. Palearctic. Nordic distribution: Sweden, Norway, Denmark, Finland. Swedish distribution: Sk, Go, Sm, Öl, Ög, Sö, Up.

Remarks. Feeding and breeding on mainly *Galium*, but also found on other plants.

Belothrips ferrugineus (Uzel, 1895)

Distribution. Palearctic. Nordic distribution: Sweden, Finland. Swedish distribution: Sö, Vr.

Remarks. First record for Sö. Feeding and breeding on Rubiaceae, here also collected on *Juncus* (Cyperaceae).

Material examined. SWEDEN • 1♀; Sö, Flens kommun, Sparreholms ekhagar nature reserve, 59.086217°N, 16.830228°E, at lake side on *Juncus*; 2023.vi.07; E. Wahlberg leg.; voucher ID: CT4T.

Belothrips morio O. M. Reuter, 1899

Distribution. Palearctic. Nordic distribution: Sweden, Finland. Swedish distribution: Sk, Sö, Up, Vr, Vb.

Remarks. First records for Sk and Vb. In flowers of many different plants, herbaceous and coniferous.

Material examined. SWEDEN • 1♀ Sk., Kristianstads kommun, Lyngsjö nature reserve, 55.932709°N, 14.067986°E, marsh with *Plantago*, *Carex*, *Dactylorhiza* and *Ranunculus*, 2024.v.21; E. Wahlberg leg.; voucher ID: FG2T. • 2♀♀; Vb, Umeå kommun, Holmsund, 63.70613°N, 20.35276°E, on Fabaceae; 2023.vi.18; E. Wahlberg leg.; voucher IDs: DI2T, DI4T.

Belothrips silvarum (Priesner, 1920)

Distribution. Palearctic. Nordic distribution: Sweden, Norway, Denmark, Finland. Swedish distribution: Sk.

Remarks. Feeding and breeding on Rubiaceae, particularly *Galium verum*.

Belothrips sordidus (Uzel, 1895)

Distribution. Palearctic. Nordic distribution: Sweden. Swedish distribution: Sö.

Remarks. First record for Sweden. Living and breeding on bedstraw (*Galium*, Rubiaceae), particularly on *Galium mollugo* and *G. lucidum*.

Material examined. SWEDEN • 1♀; Sö, Nyköpings kommun, Tovetorp research station, Skrivhuset 1, 58.94753°N, 17.14682°E, from *Achnatherum* sp. (Poaceae); 2024.viii.30; M. Ulitzka leg.; collection number: MU-S-11/1.

Belothrips validus (Karny, 1910)

Distribution. Palearctic. Nordic distribution: Sweden. Swedish distribution: Sk, Sö, Vr.

Remarks. First record for Sk. Feeding and breeding on *Galium* spp. (Rubiaceae).

Material examined. SWEDEN • 4♀♀; Sk., Kristianstads kommun, Åhus, Helge river mouth, 55.925326°N, 14.323345°E, mixed herbaceous vegetation; 2024.v.20; E. Wahlberg leg.; voucher IDs: FB8T, FC1T, FC2T, FC3T.

Bolacothrips jordani (Uzel, 1895)

Distribution. Palearctic. Nordic distribution: Norway, Finland, Denmark, Sweden. Swedish distribution: Sö.

Remarks. First record for Sweden. Found on grasses (Poaceae).

Material examined. SWEDEN • 7♀♀; Sö, Nyköpings kommun, Tovetorp research station, Skrivhuset 1, 58.94753°N, 17.14682°E, from *Achnatherum* sp. (Poaceae); 2024.viii.30; M. Ulitzka leg.; collection numbers: MU-S-10/1–7 and 2 females (apterous form) MU-S-11/5 & 6.

Bradinothrips musae (Hood, 1956)

Distribution. Nearctic. Nordic distribution: Sweden. Swedish distribution: only sporadically indoors/in greenhouses

Remarks. Breeding on leaves and feeding on fruit of *Musa*.

Ceratothrips ericae (Haliday, 1836)

Distribution. Palearctic. Nordic distribution: Sweden, Norway, Denmark, Finland, Iceland. Swedish distribution: Sk, Ha, Sm, Öl, Go, Ds, Vr, Sö, Up, Dr, Me, Vb, Ås, Pi, To.

Remarks. First records for Ås and Pi. Feeding and breeding on flowers of Ericaceae, here also collected on *Astragalus* (Fabaceae).

Material examined. SWEDEN • 1♀; Ås, Dorotea kommun, Västra Ormsjö, 64.467878°N, 15.951629°E, on *Astragalus alpinus*; 2023.vi.16; E. Wahlberg leg.; voucher ID: BU7T. • 4♀♀; Pi, Arjeplogs kommun, Mattureälven-Båldakatj, 66.297221°N, 18.327394°E, on *Calluna vulgaris*; 2023.vi.17; E. Wahlberg leg.; voucher IDs: CJ2T, CJ3T, CJ7T, CJ8T.

Chaetanaphothrips orchidii (Moulton, 1907)

Distribution. Widespread in tropical regions. Nordic distribution: Sweden, Norway, Denmark, Finland (only indoors/in greenhouses). Swedish distribution: sporadically indoors/in greenhouses.

Remarks. Feeding and breeding on leaves of different plants.

Chirothrips ambulans Bagnall, 1932

Distribution. Palearctic. Nordic distribution: Sweden, Finland. Swedish distribution: Me, Sö.

Remarks. First record for Sweden. Feeding and breeding on Poaceae, here also collected on *Juncus* (Juncaceae). Species delimitation using distance, ASAP (Fig. 11), support the validity of *C. ambulans*. Even though the tree-based species delimitation shows some ambiguity for *Chirothrips* regarding specimens within *C. manicatus*, it does also support *C. ambulans* (Fig. 12). It is also morphologically distinct from *C. manicatus* with tergite VII campaniform sensilla situated anterior to subbasal line as also used in zur Strassen (2003).

Material examined. SWEDEN • 1♀; Me, Timrå kommun, Indalsälvens delta, 62.508667°N, 17.448164°E, in mixed herbaceous vegetation; 2023.vi.14; E. Wahlberg, leg.; voucher IDs: DI7T. • 2♂♂; Sö, Nyköpings kommun, Tovetorp research station, 58.94773°N, 17.14238°E, on *Juncus*; 2024.viii.27; E. Wahlberg leg.; voucher IDs: DE5T, DE6T.

Chirothrips hamatus Trybom, 1895

Distribution. Holarctic. Nordic distribution: Sweden, Norway, Denmark, Finland. Swedish distribution: Sk, Sm, Go, Ög, Bo, Sö, Up, Vr, Jä, Ån, Ly.

Remarks. First records for Sm, Sö, Jä, Ån, and Ly. Feeding and breeding on Poaceae, pest on different grasses that affect seed viability. Prefers humid habitats.



Figure 11. Species delimitation using distance, ASAP, of *Chirothrips*. Partition with best fit is outlined with red (distance: 0.0404, p-value: 1.602413e-01, ASAP score: 1.000000, rank: 1).



Figure 12. Tree-based species delimitation with mPTP of *Chirothrips*.

Material examined. SWEDEN • 1♀; Sm, Västerviks kommun, Tindered, 57.9766°N, 16.4839°E, mixed herbaceous vegetation; 2024.vi.30; E. Wahlberg leg.; voucher ID: DX1T. • 1♀; Sö., Flens kommun, Sparreholm, beach, 59.086913°N, 16.829442°E, on *Phragmites australis*, 2024.vii.29; E. Wahlberg leg.; voucher ID: EY1T. • 7♀♀; Jä, Strömsunds kommun, Sävselet, 63.538795°N, 15.444725°E, mixed herbaceous vegetation; 2023.vi.15; E. Wahlberg leg.; voucher IDs: BY2T, BY5T, BY6T, BY7T, BY8T, BZ1T, BZ3T. • 1♀; Jä, Ragunda kommun, Krångede, 63.146760°N, 16.057726°E, roadside with herbaceous vegetation; 2023.vi.15; E. Wahlberg leg.; voucher ID: CC1T. • 5♀♀; Ån, Nordmalings kommun, Torrböle, 63.700503°N, 19.604039°E, mixed herbaceous vegetation; 2023.vi.19; E. Wahlberg leg.; CY5T, CY6T, CZ1T, CZ4T, CZ9T. • 5♀♀; Ly, Sorsele kommun, Saxnäs, 65.455847°N, 17.572265°E, mixed herbaceous vegetation; 2023.vi.16; E. Wahlberg leg.; voucher IDs: CJ9T, CK1T, CK2T, CK3T, CK4T, CK5T, CK6T.

Chirothrips manicatus Haliday, 1836

Distribution. Holarctic, introduced in Australasia. Nordic distribution: Sweden, Norway, Denmark, Finland. Swedish distribution: Sk, Bl, Ha, Sm, Öl, Go, Ög, Vg, Bo, Sö, Up, Vr, Dr, Hs, Me, Jä, Vb, Ås, Ly, Pi, Lu.

Remarks. First records for Bl, Vg, Up, Me, Jä, Ås, Ly, and Pi. Feeding and breeding on Poaceae, pest on different grasses that affect seed viability. Here also collected on *Barbarea vulgaris* (Brassicaceae), *Anchusa officinalis* (Boraginaceae) and *Astragalus alpinus* (Fabaceae).

Material examined. SWEDEN • 10♀♀; Bl, Karlskrona kommun, Knösö, 56.167848°N, 15.664502°E, on Poaceae; 2023.v.18; E. Wahlberg leg.; voucher IDs: CV9T, CW1T, CW2T, CW3T, CW4T, CW5T, CW6T, CW7T, CW8T, CW9T. • 4♀♀; Bl, Karlskrona kommun, Sandhamn, 56.090980°N, 15.856533°E, mixed herba-

ceous vegetation; 2023.v.31 E. Wahlberg leg.; voucher IDs: CQ1T, CQ2T, CQ5T, CQ6T. • 2♀♀; Bl, Karlskrona kommun, Torhamn, 56.092591°N, 15.853811°E; on *Barbarea vulgaris*; 2023.v.31; E. Wahlberg leg.; voucher IDs: CO7T, CP5T. • 6♀♀; Bl, Olofströms kommun, Kräftemåla, 56.395876°N, 14.664588°E, grazed meadow; 2023.v.31; E. Wahlberg; voucher IDs: CR7T, CR8T, CR9T, CS1T, CS2T, CS5T. • 5♀♀; Bl, Karlskrona kommun, Augerum, 56.217489°N, 15.673053°E, on Poaceae; 2023.v.30; E. Wahlberg leg.; voucher IDs: CX1T, CX2T, CX3T, CY1T, CY2T. • 1♀; Bl, Karlskrona kommun, Augerum, 56.216929°N, 15.676876°E, on *Anchusa officinalis*; 2023.vi.01; E. Wahlberg leg.; voucher ID: CM8T. • 1♀; Vg, Tjörns kommun, Tolleby, 57.97712°N, 11.60566°E, mixed herbaceous vegetation; 2022.vi.11; E. Wahlberg leg.; voucher ID: DJ8T. • 1♀; Up, Uppsala kommun, Lågbol, 60.034883°N, 18.420316°E, on *Calamagrostis*; 2021.vii.19;. E. Wahlberg leg.; voucher ID: AP3T. • 1♀; SE, Me., Timrå kommun, Indalsälvens delta, 62.508667°N, 17.448164°E, mixed herbaceous vegetation; 2023.vi.14; E. Wahlberg leg.; voucher ID: DI8T. • 3♀♀; Jä, Ragunda kommun, Boberg, 63.359057°N, 15.717947°E, on *Astragalus alpinus*; 2023.vi.15; E. Wahlberg leg.; voucher IDs: BX2T, BX3T, BX8T. • 3♀♀; Jä, Ragunda kommun, Krångede, 63.146760°N, 16.057726°E, roadside with herbaceous vegetation; 2023.vi.15; E. Wahlberg leg.; voucher IDs: CB5T, CB8T, CC2T. • 1♀; Ås, Dorotea kommun, Häggås, 64.402187°N, 16.434090°E, mixed herbaceous vegetation; 2023.vi.16; leg. E. Wahlberg leg.; voucher ID: BV9T. • 4♀♀; Ås, Dorotea kommun, Västra Ormsjö, 64.467878°N, 15.951629°E, on *Astragalus alpinus*; 2023.vi.16; E. Wahlberg leg.; voucher IDs: BV1T, BV2T, BV3T, BV5T. • 4♀♀; Ly, Storuman kommun, Stensele, 65.100466°N, 17.167255°E, mixed herbaceous vegetation; 2023.vi.16; E. Wahlberg leg.; voucher IDs: CK7T, CK8T, CK9T, CL1T. • 4♀♀; Pi, Arjeplogs kommun, Arjeplog, 66.040783°N, 17.868661°E, on *Astragalus alpinus*; 2023.vi.18; E. Wahlberg leg.; CL3T, CL4T, CL5T, CL6T.

Ctenothrips distinctus (Uzel, 1895)

Distribution. Palearctic. Nordic distribution: Sweden, Norway, Denmak, Finland. Swedish distribution: Sk, Sö, Up.

Remarks. Feeding and breeding on *Convallaria majalis*.

Dictyothrips betae Uzel, 1895

Distribution. Palearctic. Nordic distribution: Sweden, Norway, Denmark, Finland. Swedish distribution: Sk.

Remarks. Feeding and breeding on different herbaceous plants.

Drepanothrips reuteri Uzel, 1895

Distribution. Palearctic, introduced in the Neotropics. Nordic distribution: Sweden, Norway, Denmark, Finland. Swedish distribution: Vr.

Remarks. Feeding and breeding on leaves of different plants and trees, pest on *Vitis vinifera*.

Echinothrips americanus Morgan, 1913

Distribution. Nearctic, introduced worldwide. In temperate areas indoors and greenhouses. Nordic distribution: Sweden, Norway, Finland (only indoors/in greenhouses). Swedish distribution: sporadically indoors/in greenhouses.

Remarks. Feeding and breeding on leaves of many different plants.

Firmothrips firmus (Uzel, 1895)

Distribution. Palearctic. Nordic distribution: Sweden, Norway, Finland. Swedish distribution: Sk, Vr, Ån, Vb.

Remarks. First record for Ån and Vb. Feeding and breeding on Fabaceae, here also collected on meadow inflorescence and *Achillea millefolium* (Asteraceae).

Material examined. SWEDEN • 1♀; Ån, Kramfors kommun, Västhammar, 63.03017°N, 17.77373°E, mixed herbaceous vegetation; 2023.vi.19; E. Wahlberg leg.; voucher ID: DM1T. • 1♀; Ån., Nordmalings kommun, Torrböle, 63.700503°N, 19.604039°E, mixed herbaceous vegetation; 2023.vi.19; E. Wahlberg leg.; voucher ID: CZ7T. • 9♀♀; Vb, Robertsfors kommun, Rickleå, 64.101214°N, 20.929401°E, meadow with mixed flowering vegetation; 2023.vi.18; E. Wahlberg leg.; voucher IDs: CH8T, CI3T, CI5T, DQ9T, DR1T, DR2T, DR3T, DR5T, DF8T. • 2♀♀; Vb, Robertsfors kommun, Rickleå, 64.101214°N, 20.929401°E, on *Achillea millefolium*, 2023.vi.18; E. Wahlberg leg.; voucher IDs: CG3T, CG4T.

Frankliniella intonsa (Trybom, 1895)

Distribution. Palearctic but introduced worldwide. Nordic distribution: Sweden, Norway, Finland, Denmark. Swedish distribution: Sk, Bl, Ha, Sm, Öl, Go, Ög, Sö, Up, Vs, Vr, Dr, Gä, Me, Ån, Vb.

Remarks. First records for Bl, Ha, Gä, Me, and Ån. Breeding and feeding on leaves and in flowers of many different plants.

Material examined. SWEDEN • 8♀♀; Bl, Karlskrona kommun, Augerum, 56.216929°N, 15.676876°E, on *Anchusa officinalis*; 2023.vi.01; E. Wahlberg leg.; voucher IDs: CM2T, CM7T, CN3T, CN4T, CN5T, CN6T, CO3T, CP1T. • 1♀; Ha, Halmstad kommun, Särdal, beach meadow; 56.73382°N, 12.64278°E; 2021.v.19; E. Wahlberg leg.; voucher ID: AI1T. • 1♀; Gä, Gävle kommun, Östra Sikvik, 60.66559°N, 17.29876°E, mixed herbaceous vegetation; 2023.vi.20; E. Wahlberg leg.; voucher ID: DL5T. • 2♀♀; Me, Timrå kommun, Indalsälvens delta, 62.508667°N, 17.448164°E, mixed herbaceous vegetation; 2023.vi.14; E. Wahlberg leg.; voucher IDs: DJ1T, DJ2T. • 1♀; Ån, Kramfors kommun, Västhammar, 63.03017°N, 17.77373°E, mixed herbaceous vegetation; 2023.vi.19; E. Wahlberg leg.; voucher ID: DM3T. • 1♀; Ån, Nordmalings kommun, Torrböle, 63.700503°N, 19.604039°E, mixed herbaceous vegetation; 2023.vi.19; E. Wahlberg leg., voucher ID: CZ6T.

Frankliniella occidentalis (Pergande, 1895)

Distribution. Nearctic, but introduced worldwide. In temperate areas indoors and greenhouses. Nordic distribution: Sweden, Norway, Denmark, Finland (only indoors/in greenhouses). Swedish distribution: Sk.

Remarks. Breeding and feeding on leaves, flowers, and buds of many different plants.

Frankliniella pallida (Uzel, 1895)

Distribution. Palearctic. Nordic distribution: Sweden, Norway, Denmark, Finland. Swedish distribution: Sk, Sm, Up, Vs, Hs.

Remarks. Feeding and breeding on leaves and flowers of many different plants. This species prefers warmer and drier habitats.

Frankliniella tenuicornis (Uzel, 1895)

Distribution. Holarctic. Nordic distribution: Sweden, Norway, Denmark, Finland. Swedish distribution: Sk, Sm, Öl, Go, Ög, Bo, Ds, Sö, Up, Vr, Hr, Vb, Nb, Ly.

Remarks. Feeding on a variety of plants, breeding on Poaceae.

Iridothrips iridis (Watson, 1924)

Distribution. Palearctic, introduced in the Nearctic. Nordic distribution: Sweden, Norway, Denmark, Finland. Swedish distribution: Sk, Sö.

Remarks. New record for Sö. Feeding and breeding on leaves of *Iris pseudacorus*. This species prefers humid habitats.

Material examined. Sweden • 2♀♀3♂♂; Sö., Gnesta kommun, Björnlunda, 59.06726°N, 17.15507°E, on *Iris pseudacorus*; 2024.vi.03; E. Wahlberg leg.; voucher IDs: EFB5, EFB6, EFB2, EFB3, EFB4. • 2♀♀; Sö., Nyköpings kommun, Torpesta kvarn, 58.931712°N, 17.177325°E, on *Iris pseudacorus*; 2024.viii.07; E. Wahlberg leg.; voucher IDs: EM5T, EM7T.

Kakothrips pisivorus (Westwood, 1880)

Distribution. Palearctic. Nordic distribution: Sweden, Norway, Denmark, Finland. Swedish distribution: Sk, Ög, Bo, Sö, Up, Hs.

Remarks. Feeding and breeding on different species of Fabaceae.

Kakothrips priesnerorum Bournier, 1971

Distribution. Palearctic. Nordic distribution: Sweden. Swedish distribution: Vb.

Remarks. First record for Sweden. Feeding and breeding in flowers of herbaceous plants.

Material examined. SWEDEN • 1♀; Vb, Robertsfors kommun, Rickleå, 64.101214°N, 20.929401°E, meadow with mixed flowering herbaceous vegetation; 2023.vi.18; E. Wahlberg leg.; voucher ID: CH6T.

Limothrips angulicornis Jablonowski, 1894

Distribution. Palearctic, introduced worldwide. Nordic distribution: Sweden, Finland. Swedish distribution: Up.

Remarks. Feeding and breeding on leaves of Poaceae, but also occurring on other herbaceous plants and deciduous trees.

Limothrips cerealium Haliday, 1836

Distribution. Palearctic, introduced worldwide. Nordic distribution: Sweden, Norway, Denmark, Finland. Swedish distribution: Sk, Bl, Ha, Vg, Sm, Öl, Go, Ög, Bo, Sö, Up, Dr, Vb, Nb, To.

Remarks. First records for Vg and Nb. Feeding and breeding on leaves and flowers of Poaceae.

Material examined. SWEDEN • 4♀♀; Vg., Vara kommun, 58.256°N, 12.983°E, on patio in urban environment; 2025.vii.07; via Swedish Board of Agriculture in Skara. • 1♀; Nb., Luleå kommun, Gammelstad kyrkby, 65.646111°N, 22.234167°E, on *Consolida regalis*, 2025.viii.08, leg. C.-A. Gertsson.

Limothrips consimilis Priesner, 1926

Distribution. Palearctic. Nordic distribution: Sweden. Swedish distribution: Sö.

Remarks. First record for Sweden. Feeding and breeding on leaves and flowers of Poaceae in dry and warm habitats.

Material examined. SWEDEN • 4♀♀; Sö, Gnesta kommun, Önnersta, 59.0519°N, 17.1437°E, on Poaceae; 2023.vii.18; E. Wahlberg leg.; voucher IDs: BT1T, BT3T, DP8T, DP9T.

Limothrips denticornis Haliday, 1836

Distribution. Palearctic, introduced worldwide. Nordic distribution: Sweden, Norway, Denmark, Finland. Swedish distribution: Sk, Bl, Sm, Öl, Go, Ög, Vg, Bo, Nä, Sö, Up, Vs, Vr, Jä.

Remarks. First record for Bl. Feeding and breeding on leaves and flowers of Poaceae.

Material examined. SWEDEN • 2♀♀; Bl, Karlskrona kommun, Augerum, 56.217489°N, 15.673053°E, on Poaceae; 2023.v.30; E. Wahlberg leg.; voucher IDs: BI8T, CX7T.

Mycterothrips consociatus (Targioni-Tozzetti, 1886)

Distribution. Palearctic. Nordic distribution: Sweden, Finland. Swedish distribution: Sö, Vr, Ds, Up, Dr.

Remarks. First record for Sö. Feeding and breeding on the leaves of different deciduous trees, particularly on *Alnus*, *Betula*, *Corylus*, *Quercus* and *Salix*.

Material examined. SWEDEN • 1♀; Sö., Nyköping kommun, Pumptorp, roadside meadow, 58.667763°N, 16.811182°E; 2021.vi.03; E. Wahlberg leg.; voucher ID: FN9T.

Mycterothrips latus (Bagnall, 1912)

Distribution. Palearctic. Nordic distribution: Sweden, Norway, Denmark, Finland. Swedish distribution: Sk, Öl, Sö, Up, Vr, Jä, Hs, Ån, Vb, Ly, Lu, To.

Remarks. First record for Ån. Feeding and breeding on the leaves of different deciduous trees, mainly Betulaceae. Here collected on ground vegetation.

Material examined. SWEDEN • 1♀; Ån, Nordmalings kommun, Torrböle, 63.700503°N, 19.604039°E, in mixed herbaceous vegetation; 2023.vi.19; E. Wahlberg leg.; voucher ID: CY4T.

Mycterothrips salicis (O. M. Reuter, 1879)

Distribution. Holarctic. Nordic distribution: Sweden, Norway, Denmark, Finland. Swedish distribution: Sm, Öl, Go, Ög, Ds, Sö, Up, Vr, Dr, Vb, Ly.

Remarks. First record for Dr. Feeding and breeding on the leaves of different deciduous trees, mainly Salicaceae. Here collected on ground vegetation.

Material examined. SWEDEN • 3♀♀; Dr, Orsa kommun, Koppången nature reserve, 61.362499°N, 14.786772°E, on *Angelica sylvestris*; 2024.vii.30; E. Wahlberg leg.; voucher IDs: EE3T, EE5T, EE6T. • 1♀; Dr., Mora kommun, Utanmyra, 60.924194°N, 14.585461°E, in mixed herbaceous vegetation; 2024.vii.31; E. Wahlberg leg.; voucher ID: EI5T. • 1♀; Dr., Mora kommun, Utanmyra, 60.927792°N, 14.589847°E, on *Carex aquatilis*; 2024.vii.31; E. Wahlberg leg.; voucher ID: EH6T.

Odontothrips biuncus John, 1921

Distribution. Holarctic. Nordic distribution: Sweden, Norway, Denmark, Finland. Swedish distribution: Sk, Öl, Go, Vr, Ds, Up, Jä Ån, Vb, Ly, Pi, Lu, To

Remarks. First records for Up, Jä, Ån, Vb, Ly, and Pi. Feeding and breeding in flowers of Fabaceae, particularly *Vicia cracca* and *V. villosa*.

Material examined. SWEDEN • 1♀; Up, Vallentuna kommun, private garden, 59.5167°N, 18.0648°E; 2023.vi.27; T. Kronestedt leg.; voucher ID: DE8T. • 1♀; Jä, Ragunda kommun, Krångede, 63.146760°N, 16.057726°E, roadside mixed herbaceous vegetation; 2023.vi.15; E. Wahlberg leg.; voucher ID: CB9T. • 1♀1♂;

Ån, Nordmalings kommun, Torrböle, 63.700503°N, 19.604039°E, mixed herbaceous vegetation; 2023.vi.19; E. Wahlberg leg.; voucher IDs: CZ2T, CZ3T. • 3♀♀; SE, Ån., Sollefteå kommun, Junsele, 63.71736°N, 16.82341°E, mixed herbaceous vegetation; 2023.vi.19; E. Wahlberg leg.; voucher IDs: DJ9T, DK1T, DK8T. • 16♀♀2♂♂; Vb, Robertsfors kommun, Rickleå, 64.101214°N, 20.929401°E, meadow with mixed flowering herbaceous vegetation; 2023.vi.18; E. Wahlberg leg.; voucher IDs: BO4T, BO5T, BO6T, CH2T, CH5T, CH7T, CI4T, CI6T, DR4T, DR6T, DF9T, DG2T, DG3T, DG4T, DG5T, DG6T, DG7T, DG8T. • 1♀; SE, Ly., Storuman kommun, Stensele, 65.100466°N, 17.167255°E, mixed herbaceous inflorescence; 2023.vi.16; E. Wahlberg leg.; voucher ID: DQ6T. • 1♀; Pi, Arjeplogs kommun, Arjeplog, 66.040783°N, 17.868661°E, on *Astragalus alpinus*; 2023.vi.18; E. Wahlberg leg.; voucher ID: BP4T.

Odontothrips confusus Priesner, 1926

Distribution. Palearctic. Nordic distribution: Sweden. Swedish distribution: Öl.

Remarks. Feeding and breeding in flowers of Fabaceae, particularly *Medicago*.

Odontothrips intermedius (Uzel, 1895)

Distribution. Palearctic. Nordic distribution: Sweden, Norway, Denmark, Finland. Swedish distribution: Sö, Up, Vr, Me, Ly.

Remarks. First records for Me and Ly. Feeding and breeding in flowers of Fabaceae, particularly *Lathyrus*.

Material examined. SWEDEN • 2♀♀; Me, Timrå kommun, Indalsälvens delta, 62.508667°N, 17.448164°E, mixed herbaceous vegetation; 2023.vi.14; E. Wahlberg leg.; voucher IDs: DI5T, DI6T. • 2♀♀; Me, Timrå kommun, Svedje, 62.519346°N, 17.288289°E, mixed herbaceous vegetation; 2023.vi.14; E. Wahlberg leg.; voucher IDs: CC6T, CC9T. • 2♀♀; Ly, Storuman kommun, Stensele, 65.100466°N, 17.167255°E, in mixed herbaceous inflorescence; 2023.vi.16; E. Wahlberg leg.; voucher IDs: DQ7T, DQ8T.

Odontothrips loti (Haliday, 1852)

Distribution. Holarctic. Nordic distribution: Sweden, Norway, Denmark, Finland. Swedish distribution: Sk, Öl, Sö, Up, Vr, Jä, Ån, Ås, Ly.

Remarks. First records for Sö, Jä, Ån, Ås and Ly. Feeding and breeding in flowers of Fabaceae.

Material examined. SWEDEN • 1♀; Sö, Flens kommun, Malmköpings nature reserve, 59.1458°N, 16.7205°E; 2023.vi.07; E. Wahlberg leg.; voucher ID: BZ4T. • 4♀♀3♂♂; Jä, Ragunda kommun, Boberg, 63.359057°N, 15.717947°E, on *Astragalus alpinus*; 2023.vi.15; E. Wahlberg leg.; voucher IDs: BX1T, BX4T, BX5T, BX6T, BX7T, BX9T, BY1T. 1♀; Jä, Strömsunds kommun, Sävselet, 63.538795°N, 15.444725°E, mixed herbaceous inflorescences; 2023.vi.15; E. Wahlberg leg.; voucher ID: BY4T. • 1♀; Ån, Kramfors kommun, Västham-

mar, 63.03017°N, 17.77373°E, mixed herbaceous vegetation; 2023.vi.19; E. Wahlberg leg.; voucher ID: DL9T. • 2♀♀; Ån, Sollefteå kommun, Junsele, 63.71736°N, 16.82341°E, mixed herbaceous vegetation; 2023.vi.19; E. Wahlberg leg.; voucher IDs: DK4T, DK7T. • 1♀; Ås, Dorotea kommun, Västra Ormsjö, 64.467878°N, 15.951629°E, on *Astragalus alpinus*; 2023.vi.16; E. Wahlberg leg.; voucher ID: BG6T. • 1♂; Ly, Storuman kommun, Stensele, 65.100466°N, 17.167255°E, mixed herbaceous inflorescences; 2023.vi.16; E. Wahlberg leg.; voucher ID: DQ5T.

Odontothrips phaleratus (Haliday, 1836)

Distribution. Palearctic. Nordic distribution: Sweden, Norway, Denmark, Finland. Swedish distribution: Sk, Bl, Sö, Up, Vr, Dr, Ån, Me.

Remarks. First record for Ån and Me. Feeding and breeding in flowers of Fabaceae, particularly *Lathyrus pratensis* and *Vicia* spp.

Material examined. SWEDEN • 2♀♀1♂; Ån, Kramfors kommun, Västhammar, 63.03017°N, 17.77373°E, mixed herbaceous vegetation; 2023.vi.19; E. Wahlberg leg.; voucher IDs: DL8T, DM2T, DM4T. • 1♀; Me., Timrå kommun, Indalsälvens delta, 62.508667°N, 17.448164°E, in mixed herbaceous vegetation; 2023.vi.14; E. Wahlberg leg.; voucher ID: DY8T.

Odontothrips ulicis (Haliday, 1828)

Distribution. Palearctic. Nordic distribution: Sweden. Swedish distribution: Sk, Bl, Sm, Go, Ög, Bo, Up, Vs, Sö, Up.

Remarks. Feeding and breeding in flowers of *Ulex* spp. (Fabaceae).

Oxythrips ajugae Uzel, 1895

Distribution. Palearctic. Nordic distribution: Sweden, Norway, Denmark, Finland. Swedish distribution: Sk, Bl, Sm, Ög, Vg, Sö, Up, Vr, Dr, Jä, Vb, Nb, Pi, Lu, To.

Remarks. First records for Bl, Sm, Nb, and Pi. Breeding in the cones of *Pinus*, but adults found on various plants around the trees.

Material examined. SWEDEN • 1♀; Bl, Karlskrona kommun, Torhamn, 56.092591°N, 15.853811°E, on *Barbarea vulgaris*; 2023.v.31; E. Wahlberg leg.; voucher ID: CP3T. • 2♀♀; Bl, Olofströms kommun, Kräftemåla, 56.395876°N, 14.664588°E, grazed meadow; 2023.v.31; E. Wahlberg leg.; voucher IDs: CR4T, CR5T. • 1♀; Bl, Karlskrona kommun, Augerum, 56.217489°N, 15.673053°E, on Poaceae; 2023.v.30; E. Wahlberg leg.; voucher ID: CX9T. • 1♀; Sm, Nybro kommun, Alsterbo, 56.896544°N, 15.883643°E, deciduous forest with *Fraxinus*, *Quercus* and *Corylus*; 2021.v.11; E. Wahlberg leg.; voucher ID: AF1T. • 1♀; Nb, Åsele kommun, Björnlandet National Park, 63.970150°N, 18.053417°E, Malaise trap; 2011.vi.15–vi.29; K. Norberg & B.O. Johansson leg.; voucher ID: AB2T. • 1♀; Pi, Arjeplogs kommun, Mattureälven-Båldakatj, 66.297221°N, 18.327394°E, on mixed ground vegetation in old pine forest; 2023.vi.17; E. Wahlberg leg.; voucher ID: CJ5T.

Oxythrips bicolor (O. M. Reuter, 1879)

Distribution. Palearctic. Nordic distribution: Sweden, Norway, Denmark, Finland. Swedish distribution: Sk, Bl, Sm, Ög, Vg, Sö, Up, Vr, Dr, Jä, Vb, Lu, To, Pi.

Remarks. First records for Bl, Sm, Vg, and Pi. Breeding on Cupressaceae and Pinaceae, adults found feeding on a large variety of plants.

Material examined. SWEDEN • 1♀; Bl, Ronneby kommun, Värperyd, 56.295140°N, 15.290304°E, on dead *Fraxinus*; 2022.v.18; E. Wahlberg leg.; voucher ID: AX1T. • 2♀♀; Bl, Tingsryd kommun, Göljahult, 56.429934°N, 15.220758°E, on *Luzula*; 2022.v.18; E. Wahlberg leg.; voucher IDs: AX8T, AX9T. • 1♀; Bl, Tingsryd kommun, Göljahult, 56.429934°N, 15.220758°E, on *Phragmites*; 2022.v.18; E. Wahlberg leg.; voucher ID: AX6T. • 1♀; Sm., Kalmar län, Tyllinge, 58.023443°N, 16.072688°E, on *Myrica gale*; 2024.v.28; E. Wahlberg leg.; voucher ID: FE6T. • 1♀; Sm., Västerviks kommun, Hallmare, 57.866219°N, 16.754423°E, on *Bolboschoenus maritimus*; 2024.v.05; E. Wahlberg leg.; voucher ID: EB9T. • 2♀♀; Sm., Västerviks kommun, Risebo nature reserve, 58.031361°N, 16.106291°E; 2024.v.28; E. Wahlberg leg.; voucher ID: EZ7T, EZ8T. • 1♀; Vg, Laxå kommun, Finnerödja, 58.931227°, 14.346057°, window trap at bog; 2021.vi.05–vii.05; E. Wahlberg leg.; voucher ID: AM8T. • 1♀; Pi, Arjeplogs kommun, Mattureälven-Båldakatj, 66.297221°N, 18.327394°E, on mixed ground vegetation in old pine forest; 2023.vi.17; E. Wahlberg leg.; voucher ID: CJ6T.

Oxythrips ulmifoliorum (Haliday, 1836)

Distribution. Palearctic. Nordic distribution: Sweden, Norway, Finland. Swedish distribution: Go, Sö, Up.

Remarks. Feeding and breeding on leaves of Oleaceae and Ulmaceae.

Platythrips tunicatus (Haliday, 1852)

Distribution. Palearctic. Nordic distribution: Sweden, Denmark, Finland. Swedish distribution: Sö, Up.

Remarks. Feeding and breeding in flowers and on leaves of different species of *Galium*, particularly on *Galium mollugo* (Rubiaceae), but in Sweden also collected on Poaceae.

Pteridothrips pteridicola (Karny, 1914)

Distribution. Oriental. Nordic distribution: Sweden (only indoors/in greenhouses). Swedish distribution: Sk.

Remarks. Feeding and breeding on water fern (*Microsorum*).

Rhaphidothrips longistylosus Uzel, 1895

Distribution. Palearctic, introduced in the Nearctic. Nordic distribution: Sweden, Norway, Denmark, Finland. Swedish distribution: Sk.

Remarks. Feeding and breeding on Poaceae, in Sweden collected on *Orchis maculata*.

Scirtothrips longipennis (Bagnall, 1909)

Distribution. Original distribution unknown, but introduced worldwide indoors and in greenhouses. Nordic distribution: Sweden, Norway, Denmark, Finland (only indoors/in greenhouses). Swedish distribution: Ha.

Remarks. Feeding and breeding on leaves of a variety of plants, a pest on cultivated *Begonia*.

Scolothrips uzeli (Schille, 1910)

Distribution. Palearctic. Nordic distribution: Sweden, Norway. Swedish distribution: Up.

Remarks. Predator on mites on *Juniperus*.

Stenothrips graminum Uzel, 1895

Distribution. Palearctic. Nordic distribution: Sweden, Norway, Denmark, Finland. Swedish distribution: Sk, Ög, Sö, Up.

Remarks. Breeding and feeding on Poaceae.

Taeniothrips inconsequens (Uzel, 1895)

Distribution. Holarctic. Nordic distribution: Sweden, Norway, Denmark, Finland, Iceland. Swedish distribution: Sk, Ha, Öl, Go, Sm, Ög, Bo, Vg, Sö, Up, Vb, Lu.

Remarks: First records for Ha, Sm, and Vb. Feeding and breeding on different deciduous plants and trees.

Material examined. SWEDEN • 2♀♀; Ha, Falkenberg kommun, Ätrafors, 57.03428°N, 12.65945°E, deciduous forest with *Fagus*, *Quercus*, *Fraxinus*, *Malus* and *Corylus*; 2021.v.18; E. Wahlberg leg.; voucher IDs: AH1T, AH2T. • 3♀♀; Sm., Västerviks kommun, Risebo nature reserve, 58.031361°N, 16.106291°E, 2024.v.28; E. Wahlberg leg.; Voucher IDs: FA2T, FA4T, FD7T. • 1♀; Vb, Umeå kommun, Holmsund, 63.70613°N, 20.35276°E, on Fabaceae; 2023.vi.18; E. Wahlberg leg.; Voucher ID: DH9T.

Taeniothrips picipes (Zetterstedt, 1828)

Distribution. Palearctic. Nordic distribution: Sweden, Norway, Denmark, Finland, Iceland. Swedish distribution: Sk, Bl, Sm, Öl, Go, Ög, Bo, Sö, Up, Vs, Vr, Dr, Jä, Hs, Ån, Vb, Ås, Ly.

Remarks. First records for Bl, Hs, Ån, Ås, and Ly. Feeding and breeding in flowers of different plants.

Material examined. SWEDEN • 1♂; Bl, Karlskrona kommun, Sandhamn, 56.090980°N, 15.856533°E, mixed herbaceous vegetation; 2023.v.31; E. Wahlberg leg.; voucher ID: CQ4T. • 1♀; Bl, Olofströms kommun, Kräftemåla, 56.395371°N, 14.654472°E, on *Vaccinium myrtillus*; 2023.v.31; E. Wahlberg leg.; voucher ID: BL2T. • 1♀; Bl, Tingsryd kommun, Göljahult, 56.429934°N, 15.220758°E, on *Luzula*; 2022.v.18; E. Wahlberg leg.; voucher ID: AX7T. • 2♀♀; Ån, Hudiksvalls kommun, 61.747222°N, 17.129722°E, on *Melampyrum pratense*, 2025.viii.04, leg. C.-A. Gertsson • 5♀♀; Ån, Nordmalings kommun, Torrböle, 63.700503°N, 19.604039°E, mixed herbaceous vegetation; 2023.vi.19; E. Wahlberg leg.; voucher IDs: CY7T, CY9T, CZ5T, CZ8T, DA1T. • 3♀♀; Ån, Sollefteå kommun, Junsele, 63.71736°N, 16.82341°E, mixed herbaceous vegetation; 2023.vi.19; E. Wahlberg leg.; voucher IDs: DK2T, DK3T. • 2♀♀; Ås, Dorotea kommun, Båtas, 64.792075°N, 15.301708°E, on Poaceae; 2023.vi.16; E. Wahlberg leg.; voucher IDs: BR6T, BW1T. • 4♀♀; Ås, Dorotea kommun, Västra Ormsjö, 64.467878°N, 15.951629°E, on *Astragalus alpinus*; 2023.vi.16; E. Wahlberg leg.; voucher IDs: BU6T, BV4T, BV6T, BV7T. • 2♀♀; Ly, Storuman kommun, Stensele, 65.100466°N, 17.167255°E, mixed herbaceous inflorescences; 2023.vi.16; E. Wahlberg leg.; voucher IDs: DQ3T, DQ4T.

Thrips alni Uzel, 1895

Distribution. Palearctic. Nordic distribution: Sweden, Finland. Swedish distribution: Sm, Up, Vr.

Remarks. Feeding and breeding on leaves of Betulaceae and Salicaceae, but adults also found on a variety of herbaceous plants and deciduous trees.

Thrips angusticeps Uzel, 1895

Distribution. Palearctic. Nordic distribution: Sweden, Denmark, Finland. Swedish distribution: Sk, Öl, Ög, Up.

Remarks. Feeding and breeding mainly on a wide range of herbaceous plants.

Thrips atratus Haliday, 1836

Distribution. Holarctic. Nordic distribution: Sweden, Norway, Denmark, Finland, Iceland. Swedish distribution: Sk, Bl, Ha, Sm, Öl, Go, Ög, Vg, Up, Vs, Bo, Sö, Vr, Dr, Hs, Me, Hr, Ån, Vb, Nb, Ly, Lu.

Remarks. First records for Bl and Me. Feeding and breeding in variety of flowering plants, often Caryophyllaceae.

Material examined. SWEDEN • 1♀3♂♂; Bl, Karlskrona kommun, Augerum, 56.216929°N, 15.676876°E, on *Anchusa officinalis*; 2023.vi.01; E. Wahlberg leg.; voucher IDs: CL8T, CN2T, CN7T, CO2T. • 2♀♀; Me, Timrå kommun, Svedje, 62.519346°N, 17.288289°E, mixed herbaceous vegetation; 2023.vi.14; E. Wahlberg leg.; voucher IDs: CC8T, CD2T.

Thrips brevicornis Priesner, 1920

Distribution. Palearctic. Nordic distribution: Sweden, Norway, Finland. Swedish distribution: Öl.

Remarks. Feeding and breeding on a large variety of different flowering plants.

Thrips calcaratus Uzel, 1895

Distribution. Palearctic, introduced in the Nearctic. Nordic distribution: Sweden, Norway, Denmark, Finland. Swedish distribution: Sm, Ån.

Remarks. First record for Sweden. Feeding and breeding on leaves of deciduous trees, pest on *Tilia*.

Material examined. SWEDEN • 1♀; Sm., Västerviks kommun, Risebo nature reserve, 58.031361°N, 16.106291°E; 2024.v.28; E. Wahlberg leg.; voucher ID: FA5T. • 4♀♀; Ån, Kramfors kommun, Västhammar, 63.03017°N, 17.77373°E, on *Tilia cordata*; 2023.vi.19; E. Wahlberg leg.; voucher IDs: DG9T, DH1T, DH2T, DH3T.

Thrips dilatatus Uzel, 1895

Distribution. Palearctic. Nordic distribution: Sweden, Norway, Denmark, Finland. Swedish distribution: Sk, Öl, Sö, Up, Vr, Ly.

Remarks. New record for Sö. Feeding and breeding in flowers of Orobanchaceae.

Material examined. SWEDEN • 1♀; Sö., Flens kommun, Sparreholm, 59.086913°N, 16.829442°E, mixed herbaceous vegetation; 2024.vii.29; E. Wahlberg leg.; voucher ID: EFH7

Thrips discolor Haliday, 1836

Distribution. Holarctic. Nordic distribution: Sweden, Norway, Denmark, Finland. Swedish distribution: Sk, Up, Sö.

Remarks. Feeding and breeding on plants of Ranunculaceae, mainly on *Ranunculus* repens.

Thrips flavus Schrank, 1776

Distribution. Palearctic, Oriental. Nordic distribution: Sweden, Norway, Denmark, Finland. Swedish distribution: Sk, Sm, Go, Ög, Sö, Up, Vr, Dr, Hs, Ån.

Remarks. First records for Hs and Ån. Feeding and breeding in a variety of flowering plants.

Material examined. SWEDEN • 1♀; Ån, Nordmalings kommun, Torrböle, 63.700503°N, 19.604039°E, mixed herbaceous vegetation; 2023.vi.19; E. Wahlberg leg.; voucher ID: CY8T. • 1♀; Ån, Hudiksvalls kommun, 61.747222°N, 17.129722°E, on *Melampyrum pratense*, 2025.viii.04, leg. C.-A. Gertsson.

Thrips fulvipes Bagnall, 1923

Distribution. Palearctic. Nordic distribution: Sweden, Norway, Denmark. Swedish distribution: Sk.

Remarks. Feeding and breeding on *Mercurialis perennis*, adults sometimes found in flowers of other plants.

Thrips fuscipennis Haliday, 1836

Distribution. Holarctic. Nordic distribution: Sweden, Norway, Denmark, Finland. Swedish distribution: Sk, Bl, Ha, Sm, Öl, Go, Ög, Bo, Ds, Sö, Up, Vr, Dr.

Remarks. First record for Bl and Dr. Feeding and breeding in a large variety of different flowering plants.

Material examined. SWEDEN • 1♀; Bl, Karlskrona kommun, Sandhamn, 56.090980°N, 15.856533°E, mixed herbaceous vegetation; 2023.v.31; E. Wahlberg leg.; voucher ID: CQ8T. • 5♀♀; Bl, Karlskrona kommun, Torhamn, 56.092591°N, 15.853811°E, on *Barbarea vulgaris*; 2023.v.31; E. Wahlberg leg.; voucher IDs: CO8T, CO9T, CP2T, CP4T, CP6T. • 1♀; Dr., Mora kommun, Utanmyra, 60.923053°N, 14.617137°E, on Poaceae; 2024.vii.31; E. Wahlberg leg.; voucher ID: EL4T. • 2♀♀; Dr., Mora kommun, Utanmyra, 60.924194°N, 14.585461°E, in mixed herbaceous vegetation; 2024.vii.31; E. Wahlberg leg.; voucher IDs: EI3T, EI4T. • 1♀; Dr., Mora kommun, Utanmyra, 60.927792°N, 14.589847°E, on *Carex aquatilis*, 2024.vii.31; E. Wahlberg leg.; voucher ID: EI1T. • 1♀; Dr., Orsa kommun, Enåns dalgång, 61.110185°N, 14.636168°E, mixed herbaceous vegetation; 2024.vii.31; E. Wahlberg leg.; voucher ID: EJ6T.

Thrips juniperinus Linnaeus, 1758

Distribution. Palearctic. Nordic distribution: Sweden, Norway. Swedish distribution: Sö, Lu.

Remarks. Feeding and breeding on *Juniperus*.

Material examined. SWEDEN • 1♀; Sö, Nyköpings kommun, Tovetorp research station, Skrivhuset 1, 58.94753°N, 17.14682°E, from *Juniperus communis* (Cupressaceae); 2024.viii.30; M. Ulitzka leg.; collection number: MU-S-08/1.

Thrips linariae Priesner, 1927

Distribution. Palearctic. Nordic distribution: Sweden, Norway. Swedish distribution: Unknown.

Remarks. Feeding and breeding in different flowering plants. It was noted for Sweden by Kobro (2011), but regional records and physical specimens are unknown.

Thrips major Uzel, 1895

Distribution. Palearctic. Nordic distribution: Sweden, Norway, Denmark, Finland. Swedish distribution: Sk, Bl, Ha, Sm, Öl, Ög, Bo, Ds, Sö, Up, Vs, Vr, Dr, Hs, Me, Hr, Jä, Ån, Vb, Nb, Ås, Ly, Pi, Lu, To.

Remarks. First records for Me, Jä, Ås, and Pi. Feeding and breeding mainly in Rosaceae, but also found in other herbaceous plants and deciduous trees.

Material examined. SWEDEN • 1♀; Me., Timrå kommun, Indalsälvens delta, 62.508667°N, 17.448164°E, mixed herbaceous vegetation; 2023.vi.14; E. Wahlberg leg.; voucher ID: DJ3T. • 2♀♀; Me, Timrå kommun, Svedje, 62.519346°N, 17.288289°E, mixed herbaceous vegetation; 2023.vi.14; E. Wahlberg leg.; voucher IDs: CD1T, CD3T. • 1♀; Jä, Strömsunds kommun, Sävselet, 63.538795°N, 15.444725°E, mixed herbaceous inflorescences; 2023.vi.15; E. Wahlberg leg.; voucher ID: BY3T. • 1♀; Ås, Dorotea kommun, Båtas, 64.792075°N, 15.301708°E, on Poaceae; 2023.vi.16; E. Wahlberg leg.; voucher ID: BW3T. • 1♀; Ås, Dorotea kommun, Häggås, 64.402187°N, 16.434090°E, mixed herbaceous inflorescences; 2023.vi.16; E. Wahlberg leg.; voucher ID: BV8T. • 1♀; Ås, Dorotea kommun, Västra Ormsjö, 64.467878°N, 15.951629°E, on *Astragalus alpinus*; 2023.vi.16; E. Wahlberg leg.; voucher ID: BU9T. • 1♀; Pi, Arjeplogs kommun, Arjeplog, 66.040783°N, 17.868661°E, on *Astragalus alpinus*; 2023.vi.18; E. Wahlberg leg.; voucher ID: CL2T. • 4♀♀; Pi, Arjeplogs kommun, Mattureälven-Båldakatj, 66.297221°N, 18.327394°E, on mixed ground vegetation in old pine forest; 2023.vi.17; E. Wahlberg leg.; voucher IDs: CI8T, CI9T, CJ1T, CJ4T.

Thrips menyanthidis Bagnall, 1923

Distribution. Palearctic. Nordic distribution: Sweden, Norway, Finland. Swedish distribution: Vg, Go, Sö, Vr, Dr.

Remarks. First records for Vg and Sö. Feeding and breeding on *Menyanthes trifoliata*. Neither the results from the distance-based (Fig. 13) or tree-based (Fig. 14) species delimitation support this species as a synonym of *T. fuscipennis* as suggested by Pitkin (1976) and Mound et al. (1976), hence the *T. menyanthidis* is here confirmed as a valid species. Even though the campaniform sensilla on the metanotum is used as a character in e.g. Mound et al. (2018) it can be variable and dubious. The two species can be separated by postocular setae and coloration as described in the key herein.

Material examined. SWEDEN • 5♀♀; Vg, Laxå kommun, Finnerödja, 58.93127°N, 14.34668°E, bog; 2024.viii.13; E. Wahlberg leg.; voucher IDs: DU2T, DU4T. • 1♀; Sö, Nyköpings kommun, Skåraviken nature reserve, 58.81361°N, 16.73475°E, wet meadow; 2024.viii.07; E. Wahlberg leg.; voucher ID: DU7T. • 1♀; Dr., Mora kommun, Utanmyra, 60.924194°N, 14.585461°E, 2024.vii.31; E. Wahlberg leg.; voucher ID: EI8T.

Thrips minutissimus Linnaeus, 1758

Distribution. Palearctic. Nordic distribution: Sweden, Norway, Denmark, Finland. Swedish distribution Sk, Ha, Sm, Öl, Ög, Bo, Sö, Up.

Remarks. Feeding and breeding in a variety of different flowering plants and deciduous trees.

Thrips nigropilosus Uzel, 1895

Distribution. Palearctic, introduced worldwide. Nordic distribution: Sweden, Norway, Denmark, Finland. Swedish distribution: Sk, Öl, Vg, Sö, Up.

Remarks. Feeding and breeding on leaves and in flowers of different plants, pest on cultivated plants in both greenhouses and outdoors.

Thrips origani Priesner, 1926

Distribution. Palearctic, but probably introduced in many regions in the world. Nordic distribution: Sweden, Norway. Swedish distribution: Sk, Sö.

Remarks. First records for Sweden. Feeding and breeding on *Origanum*, preferably in warm and dry habitats

Material examined. SWEDEN • 2♀♀; Sk, Lunds kommun, Botaniska trädgården, 55.705833°N, 13.204167°E, on *Origanum vulgare*, 2025.viii.21, leg. C.-A. Gertsson. • 5♀♀; Sö, Nyköpings kommun, Tovetorp, 58.94782°N, 17.15014°E, on *Origanum vulgare*; 2024.ix.05; E. Wahlberg leg.; voucher IDs: DT6T, DT7T, DT8T, DT9T, DU1T.

Thrips palmi Karny, 1925

Distribution. Oriental, introduced worldwide. Nordic distribution: Sweden, Norway, Denmark, Finland (only indoors/in greenhouses). Swedish distribution: sporadically indoors/in greenhouses.

Remarks. Feeding and breeding on leaves and in flowers of large variety of different plants, pest on cultivated crops and ornamental plants.

Thrips physapus Linnaeus, 1758

Distribution. Palearctic. Nordic distribution: Sweden, Norway, Denmark, Finland. Swedish distribution: Sk, Bleking, Sm, Öl, Ög, Vg, Bo, Sö, Up, Vs, Vr, Me, Hr, Vb, Nb, Lu.

Remarks. Feeding and breeding in Asteraceae, but adults found in flowers of a variety of plants.



Figure 13. Species delimitation using distance, ASAP, of *Thrips*. *T. menyanthidis* shadowed. Partition with best fit is outlined with red (distance: 0.0100, p-value: 1.817597e-01, ASAP score: 1.000000, rank: 1).

Thrips pillichi Priesner, 1924

Distribution. Palearctic. Nordic distribution: Sweden. Swedish distribution: Bl, Jä.

Remarks. First record for Sweden. Feeding and breeding on Asteraceae.

Material examined. SWEDEN • 1♀; Bl, Olofströms kommun, Kräftemåla, 56.395876°N, 14.664588°E, grazed meadow; 2023.v.31; E. Wahlberg leg.; voucher ID: CS4T. • 1♀; Jä, Ragunda kommun, Krångede, 63.146760°N, 16.057726°E, roadside herbaceous vegetation; 2023.vi.15; E. Wahlberg leg.; voucher ID: CB7T.



Figure 14. Tree-based species delimitation with mPTP of *Thrips*. *T. menyanthidis* shadowed.

Thrips pini (Uzel, 1895)

Distribution. Palearctic. Nordic distribution: Sweden, Norway, Denmark, Finland, Iceland. Swedish distribution: Sk, Ha, Sm, Bo, Sö, Up, Vr, Dr, Jä, Vb, Nb, Lu.

Remarks. First record for Jä. Feeding and breeding on Pinaceae.

Material examined. SWEDEN • 4♀♀; SE, Jä, Undersåker kommun, Trillevallen, 63.262266°N, 13.210813°E, old growth spruce forest surrounded by wires, sweep netting twigs of spruce (*Picea*); 2023.vii.25; R. Vicente leg.; voucher IDs: BU1T, BU2T, BU3T, BU4T.

Thrips roepkei Doeksen, 1953

Distribution. Palearctic. Nordic distribution: Sweden, Norway. Swedish distribution: Sö

Remarks. First record for Sweden. Feeding and breeding on *Solanum* in humid and wet habitats

Material examined. SWEDEN • 1♀; Sö., Nyköpings kommun, Tovetorp research station, 58.9476°N, 17.1467°E; 2024.viii.27; S. Larsson leg.; voucher ID: DB2T.

Thrips simplex (Morison, 1930)

Distribution. Afrotropical, introduced worldwide. In temperate areas indoors and in greenhouses. Nordic distribution: Sweden, Norway (only indoors/in greenhouses). Swedish distribution: only sporadically indoors/in greenhouses.

Remarks. Feeding and breeding on Iridaceae, occurs as pest on *Gladiolus*. Currently only found indoors but can potentially establish in warmer climate conditions in cultivated *Gladiolus*.

Thrips tabaci Lindeman, 1889

Distribution. Palearctic, introduced worldwide. Nordic distribution: Sweden, Norway, Denmark, Finland, Iceland. Swedish distribution: Sk, Bl, Sm, Öl, Go, Ög, Vg, Bo, Sö, Up, Vs, Dr, Vr, Hr, Vb, Nb.

Remarks. First record for Bl. Feeding and breeding on a large variety of plants, sometimes also reported as a predator on mites; pest on crops of *Allium*, Apiaceae, and Brassicaceae.

Material examined. SWEDEN • 9♀♀; Bl, Karlskrona kommun, Augerum, 56.216929°N, 15.676876°E, on *Anchusa officinalis*; 2023.vi.01; E. Wahlberg leg.; voucher IDs: CL9T, CM1T, CM3T, CM4T, CM5T, CM9T, CN1T, CN9T, CO1T. • 1♀; Bl, Karlskrona kommun, Sandhamn, 56.090980°N, 15.856533°E, mixed herbaceous vegetation; 2023.v.31; E. Wahlberg leg.; voucher ID: CQ3T. • 1♀; Bl, Karlskrona kommun, Torhamn, 56.092591°N, 15.853811°E, on *Barbarea vulgaris*; 2023.v.31; E. Wahlberg leg.; voucher ID: CO6T.

Thrips trehernei Priesner, 1927

Distribution. Holarctic. Nordic distribution: Sweden, Norway, Denmark, Finland, Iceland. Swedish distribution: Sk, Bl, Sm, Öl, Ög, Sö, Up, Vr, Ly.

Remarks. First records for Bl and Ly. Feeding and breeding mainly in flowers of Asteraceae, but adults also found in flowers of a variety of different plants.

Material examined. SWEDEN • 1♀; Bl, Olofströms kommun, Kräftemåla, 56.395876°N, 14.664588°E, grazed meadow; 2023.v.31; E. Wahlberg leg.; voucher ID: CR6T. • 1♀; Ly, Storuman kommun, Stensele, 65.100466°N, 17.167255°E, mixed herbaceous inflorescences; 2023.vi.16; E. Wahlberg leg.; voucher ID: DQ2T.

Thrips urticae Fabricius, 1781

Distribution. Palearctic. Nordic distribution: Sweden. Swedish distribution: Sö.

Remarks. First record for Sweden. Feeding and breeding on *Urtica dioica*.

Material examined. SWEDEN • 5♀♀; Sö., Nyköpings kommun, Tovetorp research station, 58.94756°N, 17.14519°E, on *Urtica dioica*, 2024.ix.05, E. Wahlberg leg.; DS1T, DS2T, DS3T, DS4T, DS5T.

Thrips validus Uzel, 1895

Distribution. Holarctic. Nordic distribution: Sweden, Norway, Denmark, Finland. Swedish distribution: Sk, Bo, Öl, Ög, Sö, Up, Vs, Vr, Dr, Gä, Jä.

Remarks. First records for Gä and Jä. Feeding and breeding mainly in flowers of Asteraceae, but adults also found in flowers of a variety of different plants.

Material examined. SWEDEN • 2♀♀; Gä, Gävle kommun, Östra Sikvik, 60.66559°N, 17.29876°E, mixed herbaceous vegetation; 2023.vi.20; E. Wahlberg leg.; voucher IDs: DL4T, DL6T. • 1♀; Jä., Ragunda kommun, Krångede, 63.146760°N, 16.057726°E, roadside herbaceous vegetation; 2023.vi.15; E. Wahlberg leg.; voucher ID: CB6T.

Thrips viminalis Uzel, 1895

Distribution. Palearctic. Nordic distribution: Sweden, Norway, Finland. Swedish distribution: Bo, Up, Vr, Dr.

Remarks. Feeding and breeding on leaves of Salicaceae and Betulaceae.

Thrips vulgatissimus Haliday, 1836

Distribution. Holarctic, introduced in Australasia. Nordic distribution: Sweden, Norway, Denmark, Finland, Iceland. Swedish distribution: Sk, Öl, Ha, Sm, Ög, Vg, Bo, Sö, Up, Vs, Vr, Dr, Hs, Hr, Jä, Vb, Nb, Ås, Ly, Lu, To.

Remarks. First records for Jä, Vb, and Ås. Feeding and breeding in flowers, mainly white, of a large variety of plants.

Material examined. SWEDEN • 1♀; Jä, Strömsunds kommun, Sävselet, 63.538795°N, 15.444725°E, mixed herbaceous inflorescences; 2023.vi.15; E. Wahlberg leg.; voucher ID: BZ2T. • 1♀; Vb, Skellefteå kommun, Nilsliden, 64.944032°N, 20.367689°E, mixed herbaceous vegetation; 2023.vi.18; E. Wahlberg leg.; voucher ID: CG2T. • 1♀; Ås, Dorotea kommun, Båtas, 64.792075°N, 15.301708°E, on Poaceae; 2023.vi.16; E. Wahlberg leg.; voucher ID: BW2T.

Tmetothrips subapterus (Haliday, 1836)

Distribution. Palearctic, introduced in the Nearctic. Nordic distribution: Norway, Finland. Swedish distribution: Öl, Sö.

Remarks. First record for Sö. Feeding and breeding in flowers of Caryophyllaceae and Rubiaceae.

Material examined. SWEDEN • 1♀; Sö., Gnesta kommun, Önnersta, 59.040980°N, 17.143706°E, coniferous forest; 2021.vii.16, E. Wahlberg leg.; voucher ID: AS2T. • 1♀; Sö, Gnesta kommun, Önnersta, 59.0470°N, 17.1450°E, mixed herbaceous vegetation; 2023.vii.18; E. Wahlberg leg.; voucher ID: BS7T.

Thrips in wetland habitats

The thrips of wetlands in Sweden has not been targeted before, and knowledge of the specific diversity in the country practically non-existent. Bhatti (1998) showed that several species are hydrophilous, in contrast to the general assumption of all Thysanoptera being terrestrial, and dependent on wet biotopes including specific botanical compositions. Some species are probably understudied, with wider distribution than known due to under sampling in wet environments (Jenser 2013; Wahlberg 2023b). In Poland many of the red listed species are tied to wetland habitats (Kucharczyk and Kucharczyk 2008), four vulnerable taxa (VU) and four being near threatened (NT). As wetlands being threated environments due to e.g. exploitation and climate change (Brinson and Malvárez 2002; Kingsford et al. 2016; Graversgaard et al. 2021) it is urgent to investigate the current biodiversity. This applies in particular to species whose perhaps only habitat consists of bogs, fens, marshes and swamps with established flora and biological networks. The following thrips species, with representatives from families occurring in Sweden, are connected to wetland habitats to varying extent. The list does not include species only found indoors or in greenhouses.

Family Aeolothripidae Uzel, 1895

Rhipidothrips brunneus Williams, 1913

Found in beach and bank grasses such as *Bromus* and *Phragmites*, in Sweden only collected on grasses on sandy soil on the southern east coast (Wahlberg 2023b, 2024).

Family Phlaeothripidae Uzel, 1895

***Bolothrips dentipes* (Reuter, 1880)**

Species in this genus are often found in grasses and sedges, and *B. dentipes* is associated with wet habitats such as beach meadows and bogs. The other Swedish species in the genus, *B. bicolor* Heeger, 1852 and *B. icarus* (Uzel, 1895), are more tied to drier habitats even though *B. bicolor* has been collected in coastal areas (Wahlberg and Gertsson 2022; Wahlberg et al. 2025).

***Cephalothrips monilicornis* (Reuter, 1880)**

A very common species in many different types of grasslands, where it feeds on the tissue of various grasses and sedges. Not exclusively a wet land species, but often found together with *Bolothrips dentipes* in wetter environments (Wahlberg et al. 2025).

***Haplothrips hukkineni* Priesner, 1939**

On grasses and sedges in damp environments.

***Haplothrips statices* (Haliday, 1836)**

Feeding and breeding on *Armeria maritima* in coastal areas, but also occurring where *A. maritima* has been introduced (e.g., road banks and urban areas).

Family Thripidae

Anaphothrips badius

Tied to wet environment, often coastal where it is found on grasses and sedges, but one record here also from *Rubus* in wet area.

Baliothrips dispar

On grasses and sedges in damp environments.

Baliothrips kroli

Similar ecology as *B. dispar*, but more restricted to southern half of Sweden (Wahlberg et al. 2025).

Chirothrips hamatus

Species in this genus are common on grasses, *C. hamatus* seems particularly associated with *Alopecurus* and other grasses on shores and wetlands.

Iridothrips iridis

One of the most strikingly obligate wet land species, entirely bound to its host plant *Iris pseudacorus*. It lives within the leaf sheets protected by the plants thick fluid, only migrating to flower heads during swarming and mating (Jenser 2013). For long only known from Skåne in southern Sweden (Gertsson 2015), but recent record here also from Södermanland.

Thrips menyanthidis

As *I. iridis* an obligate wet land species, found only on *Menyanthes trifoliata*. Only recently discovered in Sweden but records show that it might be more common than previously known, at least in the southern half of the country (Wahlberg et al. 2025).

Discussion

In total, we report 98 new provincial records and ten species new to Sweden. Most provinces with the highest numbers of new records (Fig. 15) are in the north, with the exceptions of Blekinge and Södermanland. This pattern likely reflects the historical undersampling of northern Sweden; the results presented here therefore provide much-needed data for this region. Blekinge has likewise been relatively understudied. The high number of records from Södermanland largely reflects the thrips workshop held at the Tovetorp Research Station in Södermanland.

The number of recorded species in each province (Fig. 15) is in some parts a reflection of historical sampling effort, such as a high number of records in provinces around Stockholm (Södermanland and Uppland). Also, size of province is a factor. But another trend is associated with latitude. Provincial totals in our dataset are strongly structured by Sweden's main biogeographical break around the Limes Norrlandicus, which approximates the transition from boreo-nemoral/deciduous–mixed forests in the south to boreal taiga in the north (Fig. 1) and reflecting step-changes in climate, vegetation, and species ranges. It broadly follows the Lower Dalälven river corridor and, in the west, lies north of Lake Vänern; many maps place it near ~60° N, though its exact position varies locally.

Provinces south of this boundary host substantially higher numbers of known thrips than those in northern Sweden, consistent with warmer climate, longer growing seasons, and higher host-plant availability to the south. This has been demonstrated for other organism groups such as caddisflies (Gullefors 2008), beetles (Wikars 2008), and fresh water snails (von Proschwitz 2025) before.


Thripidae records per Swedish province
New records
Previously known
60
50
40
30
20
10
0
To Lu Pi Ly Ås Nb Vb Ån Jä Hr Me Hs Gä Dr Vr Vs Up Sö Nä Vg Bo Ds Ög Sm Öl Go GS Ha Bl Sk

Figure 15. Total number of known species and number of new records for each province in Sweden in the present study.

Land use and host-resource patterns likely reinforce this gradient. Roughly ~60% of Sweden's arable land occurs on the fertile southern plains with a diverse crop structure; toward the north, crop diversity decreases and arable land contracts and leys dominate the smaller cropped area (Sjulgård et al. 2022). These south–north differences in cropping systems and open habitats (field margins, flowering weeds/forbs) plausibly support more host associations for phytophagous insects in southern provinces (Haddad et al. 2001). The ecology of thrips themselves fits these patterns: most species are plant-associated, many with broad tolerance for warm, disturbed, or open habitats typical of southern agricultural landscapes (Mound 2005); thus higher plant and crop diversity should scale up to higher thrips diversity.

The wetland fauna as noted herein likely reflects a combination of host-plant availability (reed/grass dominance), brackishness/hydrology, management history, and sampling method/effort. Across Sweden, many of the thrips contributing to the "wetland fauna" are grass-associated species that breed on emergent or periodically inundated graminoids in fens, marshes, and coastal reed belts. Classic grass specialists such as *Chirothrips* spp. develop and pupate inside individual grass florets, often on stands of *Phragmites australis* and other Poaceae, a life history that concentrates populations in flowering reedbeds and sedge/grass swards when inflorescences are available. Likewise, widespread "grass thrips" such as *Anaphothrips obscurus* exploit Poaceae broadly and can occur in wet meadow/reed habitats where grasses dominate. This ecology makes reedbeds and wet grasslands disproportionately important host reservoirs for wetland thrips.

Coastal wetlands along the Baltic add further structure via brackish salinity and water-level gradients, producing mosaics of brackish reed beds and coastal/salt-influenced grasslands (Dijkema 1990); these gradients are well known to shape plant communities (Čížková et al. 2023) and therefore the host template for wetland insects. In the Nordic context, these habitats are dynamic and historically managed (grazing, mowing, reed harvesting), with management choices affecting stand age structure, patchiness, and thus invertebrate bio-

mass versus diversity, younger reeds often supporting higher biomass, older stands higher richness, arguing for mosaic management where conservation is a goal (Andersen et al. 2021).

The knowledge of the Thripidae fauna of Sweden has improved considerably but there are still missing pieces. This can be highlighted by the fact that a total of 127 species has been found in the Nordic countries. Seven of the species in Sweden are not found in any other Nordic country. The majority of species are native, but nearly 20% have probably been transported to Sweden with imported plant material.

Together, this study and earlier work on Phlaeothripidae, Aeolothripidae, and Melanthripidae have yielded numerous new provincial records and additional species new to Sweden, highlighting substantial previous gaps in both the taxonomic composition and geographic distribution of the Swedish thrips fauna. Collectively, these data refine current knowledge of thrips diversity in Sweden and, more broadly, contribute a substantial set of molecular barcodes and associated biological information.

Acknowledgements

We express our gratitude to colleagues and staff at the Lund Museum of Zoology (Sweden), the Swedish Museum of Natural History (Sweden) and the Forschungsinstitut und Naturmuseum Senckenberg (Germany), who provided valuable specimens and support. The Entomological Society in Stockholm kindly provided economical support to arrange the workshop “Thysanoptera (Insecta) – Diversity, collection, preparation, and identification”.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Use of AI

No use of AI was reported.

Funding

This project was financially supported by the Swedish Taxonomy Initiative (SLU Artdatabanken, Sweden), grant number SLU.dha.2023.4.3-247.

Author contributions

EW conceived, planned, secured funding and carried out the study, performed most laboratory and identification work as well as manuscript preparation. CAG performed additional collection and manuscript writing. MRU performed additional collection, photography, and identification, and also made substantial contributions to the key. EF did additional laboratory work and slide preparations. EN and JS provided additional collected specimens and helped in the manuscript preparation.

Author ORCIDs

Emma Wahlberg https://orcid.org/0000-0002-3732-8849
Manfred R. Ulitzka https://orcid.org/0000-0002-2639-4867
Jonas Sandström https://orcid.org/0000-0001-7100-6774

Data availability

All data supporting the findings of this study are available in the main text or Supplementary Information; the BOLD barcode dataset can be accessed at https://dx.doi.org/10.5883/DS-SETHY25.

References

Ahlberg O (1926) Svensk Insektfauna 6. Almqvist & Wiksells Boktryckeri AB, Uppsala, Sweden, 62 pp.

Andersen LH, Nummi P, Rafn J, Frederiksen CMS, Kristjansen MP, Lauridsen TL, Trøjelsgaard K, Pertoldi C, Bruhn D, Bahrndorff S (2021) Can reed harvest be used as a management strategy for improving invertebrate biomass and diversity? Journal of Environmental Management 300: 113637. https://doi.org/10.1016/j.jenvman.2021.113637

Benson DA, Cavanaugh M, Clark K, Karsch-Mizrachi I, Lipman DJ, Ostell J, Sayers EW (2012) GenBank. Nucleic Acids Research 41: D36–D42. https://doi.org/10.1093/nar/gks1195

Bhatti JS (1998) Thrips in wetland habitats. Oriental Insects 32: 363–380. https://doi.org/10.1080/00305316.1998.10433782

Brinson MM, Malvárez AI (2002) Temperate freshwater wetlands: Types, status, and threats. Environmental Conservation 29: 115–133. https://doi.org/10.1017/S0376892902000085

Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL (2009) BLAST+: Architecture and applications. BMC Bioinformatics 10: 421. https://doi.org/10.1186/1471-2105-10-421

Čížková H, Kučera T, Poulin B, Květ J (2023) Ecological basis of ecosystem services and management of wetlands dominated by common reed (*Phragmites australis*): European perspective. Diversity 15(5): 629. https://doi.org/10.3390/d15050629

Criscuolo A, Gribaldo S (2010) BMGE (Block Mapping and Gathering with Entropy): A new software for selection of phylogenetic informative regions from multiple sequence alignments. BMC Evolutionary Biology 10: 210. https://doi.org/10.1186/1471-2148-10-210

Dijkema KS (1990) Salt and brackish marshes around the Baltic Sea and adjacent parts of the North Sea: Their vegetation and management. Biological Conservation 51: 191–209. https://doi.org/10.1016/0006-3207(90)90151-E

Emerson BC (2025) Delimiting species—Prospects and challenges for DNA barcoding. Molecular Ecology 34(5): e17677. https://doi.org/10.1111/mec.17677

Ewels PA, Peltzer A, Fillinger S, Patel H, Alneberg J, Wilm A, Garcia MU, Di Tommaso P, Nahnsen S (2020) The nf-core framework for community-curated bioinformatics pipelines. Nature Biotechnology 38: 276–278. https://doi.org/10.1038/s41587-020-0439-x

Folmer O, Hoeh WR, Black MB, Vrijenhoek RC (1994) Conserved primers for PCR amplification of mitochondrial DNA from different invertebrate phyla. Molecular Marine Biology and Biotechnology 3: 294–299.

Gertsson C-A (2015) An annotated checklist of Thysanoptera (thrips) from the Nordic countries. Entomologisk Tidskrift 136: 185–198.

Gertsson C-A (2019) Nya tripsarter för Norrland. Skörvnöpparn: 47–51.

Gertsson C-A (2021) Two new Thrips species for Sweden *Haplothrips alpicola* Priesner, 1950 and *Haplothrips utae* Klimt, 1970 (Thysanoptera) along with new province records. Entomologisk Tidskrift 142: 21–30.

Gertsson C-A (2022) Tripsen *Aelothrips propinquus* (Bagnall, 1924) (Thysanoptera) ny för Öland och Bohuslän. Lucanus 27: 49–51.

Gertsson C-A, Fägerström C (2017) Två nya tripsarter (Thysanoptera) för Sverige samt tre nya landskapsfynd från Öland. Entomologisk Tidskrift 138: 131–136.

Gertsson C-A, Fägerström C, Sjödahl M (2022) Två för Skandinavien nya tripsarter (Thysanoptera): *Hoplothrips caespitis* (Uzel, 1895) och *Megalothrips bonannii* Uzel, 1895 samt nya provinsfynd. Entomologisk Tidskrift 143: 17–24.

Glenn TC, Nilsen RA, Kieran TJ, Sanders JG, Bayona-Vásquez NJ, Finger JW, Pierson TW, Bentley KE, Hoffberg SL, Louha S, others (2019) Adapterama I: Universal stubs and primers for 384 unique dual-indexed or 147,456 combinatorially-indexed Illumina libraries (iTru & iNext). PeerJ 7: e7755. https://doi.org/10.7717/peerj.7755

Graversgaard M, Jacobsen BH, Hoffmann CC, Dalgaard T, Odgaard MV, Kjaergaard C, Tonderski K (2021) Policies for wetlands implementation in Denmark and Sweden–historical lessons and emerging issues. Land Use Policy 101. https://doi.org/10.1016/j.landusepol.2020.105206

Gullefors B (2008) Limes norrlandicus-a natural biogeographical border for caddisflies (Trichoptera) in Sweden. Ferrantia 55: 61–65.

Haddad NM, Tilman D, Haarstad J, Ritchie M, Knops JM (2001) Contrasting effects of plant richness and composition on insect communities: A field experiment. The American Naturalist 158: 17–35. https://doi.org/10.1086/320866

Hoang DT, Chernomor O, Von Haeseler A, Minh BQ, Vinh LS (2018) UFBoot2: Improving the ultrafast bootstrap approximation. Molecular Biology and Evolution 35: 518–522. https://doi.org/10.1093/molbev/msx281

Jenser G (2013) *Iridothrips iridis* (Watson, 1924)(Thysanoptera: Thripidae): an unusual species of thrips that lives in a subaquatic habitat. Acta Zoologica Academiae Scientiarum Hungaricae 59: 61–66. https://doi.org/10.5281/zenodo.5736083

Kalyaanamoorthy S, Minh BQ, Wong TK, Von Haeseler A, Jermiin LS (2017) ModelFinder: Fast model selection for accurate phylogenetic estimates. Nature Methods 14: 587–589. https://doi.org/10.1038/nmeth.4285

Kapli P, Lutteropp S, Zhang J, Kobert K, Pavlidis P, Stamatakis A, Flouri T (2017) Multirate Poisson tree processes for single-locus species delimitation under maximum likelihood and Markov chain Monte Carlo. Bioinformatics (Oxford, England) 33: 1630–1638. https://doi.org/10.1093/bioinformatics/btx025

Katoh K, Standley DM (2013) MAFFT multiple sequence alignment software version 7: Improvements in performance and usability. Molecular Biology and Evolution 30: 772–780. https://doi.org/10.1093/molbev/mst010

Kingsford RT, Basset A, Jackson L (2016) Wetlands: Conservation's poor cousins. Aquatic Conservation 26: 892–916. https://doi.org/10.1002/aqc.2709

Kobro S (2011) Checklist of Nordic Thysanoptera. Norwegian Journal of Entomology 58: 20–26. https://doi.org/10.5281/zenodo.16005826

Kobro S, Ultizka MR (2021) *Thrips orarius* sp. n. and six first records of thrips (Thysanoptera) from Norway. Norwegian Journal of Entomology 68: 130–145.

Kucharczyk H, Kucharczyk M (2008) The red list of threatened thrips species (Thysanoptera, Insecta) of middle-eastern Poland. Acta Phytopathologica et Entomologica Hungarica 43: 297–305. https://doi.org/10.1556/APhyt.43.2008.2.13

Mound LA (2005) Thysanoptera: Diversity and interactions. Annual Review of Entomology 50: 247–269. https://doi.org/10.1146/annurev.ento.49.061802.123318

Mound LA, Morison GD, Pitikin BR, Palmer JMT (1976) I. Part 1 Handbooks for the identification of British Insects. Royal Entomological Society, London, England, 110 pp.

Mound LA, Collins DW, Hastings A (2018) Thysanoptera Britannica et Hibernica - Thrips of the British Isles. Lucidcentral.org, Identic Pty Ltd, Queensland, Australia. https://keys.lucidcentral.org/keys/v3/british_thrips/

Nguyen LT, Schmidt HA, von Haeseler A, Bui MQ (2015) IQ-TREE: A fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. Molecular Biology and Evolution 32(1): 268–274. https://doi.org/10.1093/molbev/msu300

Pitkin B (1976) Notes on *Thrips physapus* L., *hukkineni* Priesner and *fuscipennis* Haliday (Thysanoptera: Thripidae) in Britain. Entomologist's Gazette 27: 173–178.

Puillandre N, Brouillet S, Achaz G (2021) ASAP: Assemble species by automatic partitioning. Molecular Ecology Resources 21: 609–620. https://doi.org/10.1111/1755-0998.13281

Qvick U (1977) New records and notes on the Swedish Thrips Fauna (Thysanoptera. Entomologisk Tidskrift 98: 127–130.

Ranasinghe UGSL, Eberle J, Thormann J, Bohacz C, Benjamin SP, Ahrens D (2022) Multiple species delimitation approaches with COI barcodes poorly fit each other and morphospecies–An integrative taxonomy case of Sri Lankan Sericini chafers (Coleoptera: Scarabaeidae). Ecology and Evolution 12: e8942. https://doi.org/10.1002/ece3.8942

Ratnasingham S, Hebert PD (2007) BOLD: The Barcode of Life Data System (http://www.barcodinglife.org). Molecular Ecology Notes 7: 355–364. https://doi.org/10.1111/j.1471-8286.2007.01678.x

Schliephake G, Klimt KT, Fransenflügler (1979) Die Tierwelt Deutschlands.

Sjulgård H, Colombi T, Keller T (2022) Spatiotemporal patterns of crop diversity reveal potential for diversification in Swedish agriculture. Agriculture, Ecosystems & Environment 336: 108046. https://doi.org/10.1016/j.agee.2022.108046

ThripsWiki (2026) ThripsWiki – providing information on the World's thrips. https://thrips.info/wiki/Main_Page

Vasiliu-Oromulu L, Zur Strassen R, Larsson H (2001) The systematic revision of Thysanoptera species from Swedish fauna and their geographical distribution. Entomologica Romanica 6: 93–101.

Vierstraete AR, Braeckman BP (2022) Amplicon_sorter: A tool for reference-free amplicon sorting based on sequence similarity and for building consensus sequences. Ecology and Evolution 12: e8603. https://doi.org/10.1002/ece3.8603

von Proschwitz T (2025) The Swedish Fauna of Freshwater Snails—An Overview of Zoogeography and Habitat Selection with Special Attention to Red-Listed. Species Diversity: An International Journal for Taxonomy, Systematics, Speciation, Biogeography, and Life History Research of Animals 17: 251. https://doi.org/10.3390/d17040251

Wahlberg E (2023a) A modern workflow for non-destructive DNA extraction and slide preparation of thrips (Insecta, Thysanoptera) for taxonomic studies and collection deposition. Norwegian Journal of Entomology 70: 1–5. https://doi.org/10.5281/zenodo.15883419

Wahlberg E (2023b) *Rhipidothrips brunneus* (Thysanoptera: Aeolothripidae) ny art för Sverige, samt ett nytt landskapsfynd av *Thrips menyanthidis* Bagnall, 1923 (Thripidae). Entomologisk Tidskrift 144: 47–51.

Wahlberg E (2024) The Swedish Aeolothripidae and Melanthripidae (Thysanoptera) with a Redescription of *Rhipidothrips niveipennis* Reuter, 1899. Taxonomy 4: 163–183. https://doi.org/10.3390/taxonomy4010009

Wahlberg E, Gertsson C-A (2022) Identification key to and checklist of the Swedish Phlaeothripidae (Thysanoptera). ZooKeys 1096: 161–187. https://doi.org/10.3897/zookeys.1096.83011

Wahlberg E, Gertsson C-A, Nyström D (2025) Nya tripsfynd (Thysanoptera) i den svenska faunan. Entomologisk Tidskrift 146: 97–106.

Wikars L-O (2008) Åtgärdsprogram för björklevande vedskalbaggar i Norrland. Naturvårdsverket Rapport 5843.

zur Strassen R (1959) Eight synonyms amongst the European species of Chirothrips Haliday 1836 (Thysanoptera). Journal of the Entomological Society of Southern Africa 22: 88–107.

zur Strassen R (1982) Thysanopterologische notizen (6) (Insecta: Thysanoptera). Senkenbergiana biologica 63: 191–209.

zur Strassen R (2003) Die Terebranten Thysanopteren Europas und des Mittelmeer-Gebietes. Goecke & Evers, Keltern, Germany, 277 pp.

Supplementary material 1

BOLD and GenBank accession numbers

Authors: Emma Wahlberg, Carl-Axel Gertsson, Manfred R. Ulitzka, Erika Frisk, Ellen Nein, Jonas Sandström

Data type: xlsx

Copyright notice: This dataset is made available under the Open Database License (http://opendatacommons.org/licenses/odbl/1.0/). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: https://doi.org/10.3897/zookeys.1270.180704.suppl1